

**University of South Bohemia in České Budějovice
Faculty of Science**



Silica-scaled chrysophytes of Southern Bohemian water bodies, including *Mallomonas conspersa* Dürschmidt with occurrence so far reported from Japan and New Zealand

RNDr. Thesis

Mgr. Kateřina Janatková

České Budějovice
2013

This thesis should be cited as

Janatková, K., 2013: Silica-scaled chrysophytes of Southern Bohemian water bodies, including *Mallomonas conspersa* Dürschmidt with occurrence so far reported from Japan and New Zealand. RNDr. Thesis, University of South Bohemia, Faculty of Science, České Budějovice, Czech Republic

› Annotation

We investigated the species diversity of silica-scaled chrysophytes of Borkovická Blata peat-bog and Novohradské hrady ponds using the transmission electron microscopy. We recorded 25 taxa representing five genera. In this study, we found 3 taxa, which were recorded for the first time from the territory of the Czech Republic (*Mallomonas conspersa*, *M. pumilio* var. *munda*, *Synura mollispina*). The taxon *Mallomonas conspersa* has been so far reported only from Japan and New Zealand.

› Declaration [in Czech]

Prohlašuji, že svoji rigorózní práci jsem vypracovala samostatně pouze s použitím pramenů a literatury uvedených v seznamu citované literatury.
Prohlašuji, že v souladu s § 47b zákona č. 111/1998 Sb. v platném znění souhlasím se zveřejněním své rigorózní práce, a to v úpravě vzniklé vypuštěním vyznačených částí archivovaných Přírodovědeckou fakultou elektronickou cestou ve veřejně přístupné části databáze STAG provozované Jihočeskou univerzitou v Českých Budějovicích na jejích internetových stránkách, a to se zachováním mého autorského práva k odevzdanému textu této kvalifikační práce. Souhlasím dále s tím, aby toutéž elektronickou cestou byly v souladu s uvedeným ustanovením zákona č. 111/1998 Sb. zveřejněny posudky školitele a oponentů práce i záznam o průběhu a výsledku obhajoby kvalifikační práce. Rovněž souhlasím s porovnáním textu mé kvalifikační práce s databází kvalifikačních prací Theses.cz provozovanou Národním registrem vysokoškolských kvalifikačních prací a systémem na odhalování plagiátů.

České Budějovice, 5. 11. 2013

Mgr. Kateřina Janatková

› **This thesis is based on the following publication:**


JANATKOVÁ, K. & NĚMCOVÁ, Y. (2009): Silica-scaled chrysophytes of Southern Bohemian water bodies, including *Mallomonas conspersa* Dürschmidt with occurrence so far reported from Japan and New Zealand - Fottea 9: 93-99.

IF = 1.37

› **Co-author agreement:**

The co-author fully acknowledges that Kateřina Janatková is the first author of this publication. Kateřina Janatková was responsible for field sampling and chemical analyses of the water samples. She participated in transmission electron microscopy investigations, species determination, and writing of the manuscript.

The co-author hereby consents to the publication in the RNDr. thesis of Kateřina Janatková and supports this statement with her signature:



Doc. RNDr. Yvonne Němcová, Ph.D.
Charles University in Prague
Department of Botany

› **Acknowledgement**

This study was supported by the grant No. 206/08/ P281 of the Czech Science Foundation and the Research Grant No. 21620828 of the Czech Ministry of Education.

Silica-scaled chrysophytes of Southern Bohemian water bodies, including *Mallomonas conspersa* DÜRRSCHMIDT with occurrence so far reported from Japan and New Zealand

Kateřina JANATKOVÁ & Yvonne NĚMCOVÁ*

Department of Botany, Charles University in Prague, Benátská 2, CZ-128 01 Praha 2, Czech Republic;
*corresponding author: e-mail: ynemcova@natur.cuni.cz

Abstract: Silica-scaled chrysophytes are often important components of the phytoplankton community of freshwater lakes and reservoirs. A total of 25 taxa were recorded during an investigation of Borkovická Blata peat-bog and Novohradské Hory ponds, representing five genera. Along with cosmopolitan and widely distributed species, rare *Synura mollispina*, *Mallomonas conspersa*, *M. retifera* and *M. kalinae* were found. This study also represent only the third time that *M. conspersa* has ever been reported and documented by scale microphotographs. This taxon has so far been reported only from Japan and New Zealand. The close resemblance of scale morphology to *M. retifera* is discussed. *Synura mollispina*, *M. conspersa* and *M. pumilio* var. *munda* are recorded for the first time from the territory of the Czech Republic.

Key words: Silica-scaled chrysophytes, Chrysophyceae, Synurophyceae, *Mallomonas conspersa*

Introduction

Silica-scaled chrysophytes are often important components of the phytoplankton community of freshwater lakes and reservoirs (SIVER 1995). They comprise free living and colonial flagellates from the classes Synurophyceae and Chrysophyceae (Stramenopila). Their cells are enclosed within a case formed by overlapping, species-specific, silica scales. Taxonomy within this group is still almost exclusively based on morphology of silica structures studied by a transmission or scanning electron microscopy (e.g. SIVER 1991, KRISTIANSEN 2002). Species with a narrow ecological optimum could be considered suitable bioindicators of recent and fossil environments (MUNCH 1980, SMOL 1995).

Both areas investigated for this report represent unique biotopes. Borkovická Blata peat-bog (49°14'9.133"N; 14°37'25.632"E) originated before more than ten thousand years. A thickness of the peat layer reached four to eight meters. The blocks of peat were manually dug and they were used for heating till the middle of the 20th century. Then the industrial mining started here and it finished when the main deposits were empty. The samples were collected from small shallow pools, which originated in consequence of peat exploitation, and are situated in the preserved part

of an area (54.5 ha), which has been protected since 1980. The Novohradské Hory (48°39'23.129"N; 14°40'56.122"E) represent a valuable untouched nature area with a lot of natural forests situated along Czech-Austrian border. This border area was closed to the public during the communist era (1948-1989). The Nature Park Novohradské Hory was founded in 2000. Investigated water bodies (so called "klauza", from Latin *clausura*) were established at the end of the 18th century to facilitate a wood floatation. A brief description of investigated sites is given in Table 1. The Czech Republic belongs to relatively well explored areas in central Europe (e.g. FOTT 1955, NĚMCOVÁ et al. 2003, ŘEZÁČOVÁ & NEUSTUPA 2007), nevertheless still plenty of biologically valuable water habitats have, to date, not been investigated for the presence of silica-scaled chrysophytes. The purpose of this study is to expand our knowledge of autecology and biogeography of sampled species, and to extend the list of taxa reported from the territory of the Czech Republic. This study also represent only the third time that *Mallomonas conspersa* has ever been reported and documented by scale microphotographs.

Material and Methods

Plankton net (mesh 20 μm) collections were made during October and November 2007. Water temperature, pH and conductivity were measured at the time of collection with Combo pH EC (Hanna Instruments). Concentration of ammonium ions $[\text{NH}_4^+]$, total nitrogen, total phosphorus and silica $[\text{SiO}_2]$ were obtained in a laboratory using DR/890 Colorimeter (Hach). Water samples were centrifuged or concentrated by sedimentation. Drops of the sample were dried onto formvar coated grids. Dried material was washed by repeated transfer of the grid into drops of ionized water dispensed on the hydrophobic surface of a Parafilm strip (Pechiney Packing). Dried grids were examined with a JEOL 1011 transmission electron microscope.

Results and discussion

Silica-scaled chrysophytes found during this survey, accompanied with taxonomic authorities, are listed in Table 2. A total of 25 taxa were recorded in the five investigated localities, representing five genera. The number of taxa found in a given site in Borkovická Blata peat-bog (localities 1–3) ranged from 2 to 12, Novohradské Hory ponds (localities 4–5) hosted identically 11 taxa. The main physico-chemical parameters of the investigated water bodies are summarized in Table 1. While the sampling sites of both areas were relatively similar in pH, they differed in concentration of dissolved salts and nutrient content. The most common and abundant species observed in at least four study sites included *Synura petersenii*, *S. spinosa* and *Mallomonas papillosa*. The chrysophyte flora of Southern Bohemian localities consisted mainly of pH indifferent widely distributed taxa. Along with frequently reported taxa the rare species *Synura mollispina*, *Mallomonas conspersa*, *M. retifera* and *M. kalinae* were found.

Synura mollispina represents a widely distributed but scattered species, recorded from Europe, Northern and Southern America (KRISTIANSEN & PREISIG 2007). It differs from *S. spinosa* in the presence of a secondary hexagonal reticulum that covers almost completely the scale. The scales are often elongate with slightly pointed shape (CRONBERG & KRISTIANSEN 1980), but also rounded scales were noticed (ASMUND 1968, this study, see Fig. 2).

Mallomonas conspersa was reported from Japan, as an unidentified species (TAKAHASHI

Table 1. Brief description of investigated sites accompanied with the main physico-chemical parameters and position data [(Cond.) conductivity, (Temp) temperature, (N) total nitrogen, (P) total phosphorus, (NH_4^+) ammonium ions concentration].


No.	Locality	Description	Latitude	Longitude	pH	Cond. $\mu\text{S}\cdot\text{cm}^{-1}$	Temp. $^{\circ}\text{C}$	NH_4^+ $\mu\text{g}\cdot\text{l}^{-1}$	N $\mu\text{g}\cdot\text{l}^{-1}$	P $\mu\text{g}\cdot\text{l}^{-1}$	Date of sampling
1.	Borkovická Blata peat-bog	A small artificial lake	49°14'3.971"N	14°37'36.575"E	5.6	149	6.7	2.16	2.4	0.31	19. 10. 07
2.	Borkovická Blata peat-bog	A small shallow pool	49°14'8.173"N	14°37'24.229"E	5.5	101	6.0	1.61	6.9	0.23	19. 10. 07
3.	Borkovická Blata peat-bog	A shallow pool with tufts of grass	49°14'9.133"N	14°37'25.632"E	5.9	85	6.8	1.87	2.8	0.35	19. 10. 07
4.	Novohradské Hory	A forest pond Uhlíšťský ice covered	48°38'48.07"N	14°39'19.962"E	5.6	45	0.2	0.41	1.8	0.18	14. 12. 07
5.	Novohradské Hory	A forest pond Hut'ský ice covered	48°39'23.129"N	14°40'56.122"E	5.9	46	0.0	0.21	1.7	0.2	14. 12. 07

Table 2. The complete list of all taxa found during this survey, accompanied with taxonomic authorities. Referred previous CZ records [(1) PICHRTOVÁ et al. 2007, (2) ŘEZÁČOVÁ 2006, (3) NOVÁKOVÁ et al. 2004, (4) ŘEZÁČOVÁ et al. 2004, (5) NĚMCOVÁ et al. 2003, (6) PICHRTOVÁ & VESELÁ (in press)]. Asterisk denotes a new record for the Czech Republic.

Taxon		Locality No.					Previous CZ records
		1	2	3	4	5	
<i>S. echinulata</i> KORSHIKOV	Fig. 3	X					5
<i>S. mollispina</i> (J.B.PETERSEN et J.B.HANSEN)	Fig. 2	X					*
L.S.PÉTERFI et MOMEU							
<i>S. petersenii</i> KORSHIKOV	Figs 18–19	X	X	X	X	X	1; 4
<i>S. sphagnicola</i> (KORSHIKOV) KORSHIKOV		X	X				4; 5
<i>Synura spinosa</i> KORSHIKOV		X	X		X	X	1; 4; 5
<i>S. uvella</i> EHRENBERG	Fig. 1						5
<i>M. acaroides</i> PERTY emend. IWANOFF	Fig. 15		X	X	X	X	3; 4; 5
<i>M. calceolus</i> D.E.BRADLEY					X		1; 3; 4; 5
<i>M. conspersa</i> DÜRRSCHMIDT	Figs 7–9		X				*
<i>M. costata</i> DÜRRSCHMIDT	Fig. 6		X				1; 4; 5
<i>M. crassisquama</i> (ASMUND) FOTT						X	1; 4; 5
<i>M. heterospina</i> J.W.G.LUND	Fig. 11				X	X	1; 4; 5
<i>M. kalinae</i> ŘEZÁČOVÁ	Fig. 10					X	2
<i>M. multisetigera</i> DÜRRSCHMIDT					X		6
<i>M. ouradion</i> K.HARRIS et D.E.BRADLEY						X	5
<i>M. papillosa</i> K.HARRIS et D.E.BRADLEY	Figs 13–14	X	X		X	X	3; 4; 5
emend. K.HARRIS							
<i>M. pumilio</i> var. <i>pumilio</i> K.HARRIS et D.E.BRADLEY emend. ASM., CRONB. et DÜRRS.		X	X				4; 5
<i>Mallomonas pumilio</i> var. <i>munda</i> ASMUND, CRONBERG et DÜRRSCHMIDT	Fig. 4		X		X		*
<i>M. retifera</i> DÜRRSCHMIDT	Fig. 7		X				1; 4
<i>M. schwemmleri</i> GLENK emend. GLENK et FOTT	Figs 16–17					X	1; 3; 4; 5
<i>M. striata</i> K.HARRIS et D.E.BRADLEY	Fig. 12		X		X	X	1; 3; 4; 5
<i>M. tonsurata</i> TEILING			X			X	1; 3; 4; 5
<i>Chryso-sphaerella brevispina</i> KORSHIKOV	Figs 21–22				X		5
<i>Paraphysomonas vestita</i> DE SAEDELEER	Fig. 23				X		1; 5
<i>Spiniferomonas trioralis</i> E.TAKAHASHI	Fig. 20					X	1; 5

1959) and later it was described from New Zealand (DÜRRSCHMIDT 1986). *Mallomonas conspersa* was also listed in two Japanese species surveys, but the presence was not documented by scale microphotographs (ITO 1990, 1991). *M. conspersa* was delimited from *M. calceolus* (DÜRRSCHMIDT 1986), but the structure of the scales is also very similar to that of the scales of *M. retifera* (DÜRRSCHMIDT 1982). The most distinctive difference between these two species should be the sizes of the entire cell and the scales (*M. conspersa*: cell size 8–15 x 5–7 µm, scale size: 2.8–3.5 x 1.5–2 µm; *M. retifera*: cell size 17–20 x 5–9 µm, scale size 3.2–3.7 x 2–2.4 µm), but we have revealed scales of almost the same size for both taxa (compare Figs 6 with 8 and 9). Both discussed species were found at the same locality (see Table 2) during this survey. SIVER (1991) hypothesized, on the bases of very similar scale morphology, that these two rare

taxa may represent the same species. However, scales of *M. retifera* have a transverse rib on the shield positioned just behind the dome, the character that clearly distinguishes them from those of *M. conspersa*. The posterior flanges of investigated scales were not smooth, but possessed inconspicuous struts, radiated from the V-rib (Fig. 6). These struts have not been mentioned in *M. conspersa*. As the species concept is based on scale morphology (SIVER 1991, KRISTIANSEN 2002), the border between two species may be artificial and always reflecting a personal approach of the scientists who described them. While *M. retifera* was reported from Europe (e.g. HARTMANN & STEINBERG 1989, PICHRTOVÁ et al. 2007), South America (DÜRRSCHMIDT 1980, 1982) and East Asia (ITO 1990), *M. conspersa* was thought to be an extremely rare species. This species may be widely distributed or cosmopolitan, but forming only relatively small populations compared to




Figs 1–9. Silica-scaled chrysophytes of Borkovická Blata: (1) *Synura uvella*; (2) *S. mollispina*; (3) *S. echinulata*; (4) *Mallomonas pumilio* var. *munda*; (5) *M. costata*; (6) *M. retifera*; (7–9) *M. conspersa* [(7) an irregular apical scale with a prominent dome, bearing a bristle, (8–9) body scales]. Scale bars 1 μm .

frequently recorded species. The probability to reveal the scales on electron microscopic preparations is thus much lower, despite of the considerable degree of sampling effort. FINLAY & CLARKE (1999) demonstrated positive correlation between worldwide commonness of the species, and its local abundance. *M. conspersa* was described from a small pool in a semi-swamp forest, with pH of 5.6 and conductivity $90 \mu\text{S}\cdot\text{cm}^{-1}$ (DÜRRSCHMIDT 1986). These environmental

conditions are comparable to those of recorded on Loc. 2, where this species was sampled in this study.

M. kalinae is a widespread, but rare, taxon found primarily at higher temperatures (DÜRRSCHMIDT & CROOME 1985, CROOME & TYLER 1988, ŘEZÁČOVÁ 2006). *M. kalinae* was discovered in Loc. 5 under the ice cover, where the temperature reached $0 \text{ }^{\circ}\text{C}$, thus this species has probably wider temperature tolerance. *M.*




Figs 10–17. Silica-scaled chrysophytes of Novohradské Hory: (10) *Mallomonas kalinae*; (11) *M. heterospina*; (12) *M. striata*; (13–14) *M. papillosa* [(13) body scales, (14) scales bearing bristles]; (15) *M. acaroides*; (16–17) *M. schwemmlei* [(16) a scale from a posterior part of the cell, (17) a whole scale case, scale bar 5 μm]. Other scale bars 1 μm .

pumilio has been shown to occur predominantly at temperatures less than 13 °C. There seems not to be a difference in the ecological requirements of the varieties (SIVER 1991). Both sampled varieties (*pumilio* and *munda*) have a bipolar distribution (KRISTIANSEN & PREISIG 2007). *M. pumilio* var. *munda* differs from the nominal variety in the presence of a regularly built hexagonal reticulum of the shield, with mostly six pores enclosed in each mesh. The important character is also the

presence of struts on the anterior flanges.

M. schwemmlei is best described as winter species, distributed over the Northern temperate regions. Abundant populations of *M. schwemmlei* were discovered in Hut'ský pond (Loc. 5) under the ice cover. Chrysophytes were often reported to form high population densities at low temperatures (e.g. SANDGREN 1988, SIVER 1995). They may be better adapted to surviving conditions under an ice cover (low temperature combined with



Figs 18–23. Silica-scaled chrysophytes of Novohradské Hory: (18–19) *Synura petersenii* [(18) body scales, (19) a partly disintegrated scale case]; (20) *Spiniferomonas* cf. *trioralis*, spine; (21–22) *Chrysosphaerella brevispina* [(21) a plate scale, (22) a spine scale]; (23) *Paraphysomonas vestita*, a spine scale. Scale bars 1 μ m.

low light intensities), giving them the advantage once the ice disappears (SIVER 1995). Taxa from the other genera were represented by widely distributed species *Chrysosphaerella brevispina*, *Paraphysomonas vestita* and *Spiniferomonas* cf. *trioralis*, with wide ecological tolerances. The presence of *Synura mollispina*, *Mallomonas conspersa* and *M. pumilio* var. *munda* are recorded for the first time from the territory of the Czech Republic.

Acknowledgements

This study was supported by the grant No. 206/08/P281 of the Czech Science Foundation and the Research Grant No. 21620828 of the Czech Ministry of Education.

References

- ASMUND, B. (1968): Studies on Chrysophyceae from some ponds and lakes in Alaska. VI. Occurrence of *Synura* species. – *Hydrobiologia* 31: 497–515.
- CROOME, R. & TYLER, P. (1988): Further observations of silica-scaled Chrysophyceae (Paraphysomonadaceae and Mallomonadaceae) from Australian freshwaters. – *Nova Hedwigia* 46: 481–489.
- CRONBERG, G. & KRISTIANSEN, J. (1980): Synuraceae and other Chrysophyceae from central Smaland. – *Sweden. Bot. Notiser* 133: 595–618.
- DÜRRSCHMIDT, M. (1980): Studies on the Chrysophyceae from Rio Cruces, Prov. Valdivia, South Chile by Scanning and transmission microscopy. – *Nova Hedwigia* 33: 353–388.
- DÜRRSCHMIDT, M. (1982): *Mallomonas parvula* sp. nov. and *Mallomonas retifera* sp. nov. (Chrysophyceae, Synuraceae) from South Chile. – *Can. J. Bot.* 60: 651–656.
- DÜRRSCHMIDT, M. (1986): New species of the genus *Mallomonas* (Mallomonadaceae, Chrysophyceae) from New Zealand. – In: KRISTIANSEN, J. & ANDERSEN, R.A. (eds): *Chrysophytes. Aspects and Problems*. – pp. 87–106, Cambridge University Press, Cambridge.
- DÜRRSCHMIDT, M. & CROOME, R. (1985): Mallomonadaceae (Chrysophyceae) from Malaysia and Australia. – *Nord. J. Bot.* 5: 285–298.

- FINLAY, B.J. & CLARKE, K.J. (1999): Apparent global ubiquity of species in the protist genus *Paraphysomonas*. – *Protist* 150: 419–430.
- FOTT, B. (1955): Scales of *Mallomonas* observed in the electron microscope. – *Preslia* 27: 280–282.
- HARTMANN, H. & STEINBERG, C. (1989): The occurrence of silica-scaled chrysophytes in some central European lakes and their relation to pH. – *Beih. Nov.Hedw.* 95: 131–158.
- ITO, H. (1990): Chrysophytes in the southern part of Hyogo prefecture, Japan (I). Chrysophyte flora in three ponds and reservoir. – *Jap. J. Phycol.* 38: 327–332.
- ITO, H. (1991): Chrysophytes in the southern part of Hyogo prefecture, Japan (II). *Mallomonas*. – *Jap. J. Phycol.* 39: 253–262.
- KRISTIANSEN, J. (2002): The genus *Mallomonas* (Synurophyceae) - A taxonomic survey based on the ultrastructure of silica scales and bristles. – *Opera Botanica* 139: 5–128.
- KRISTIANSEN, J. & PREISIG, H.R. (2007): Chrysophyte and Haptophyte Algae. Teil 2 / Part 2: Synurophyceae. – In: BÜDEL, B., GÄRTNER, G., KRIENITZ, L., PREISIG, H.-R. & SCHAGERL, M. (eds): Süßwasserflora von Mitteleuropa, Bd. 01/2, Freshwater Flora of Central Europe, Vol. 01/2. – 252 pp., Spektrum Akademischer Verlag.
- MUNCH, C.S. (1980): Fossil diatoms and scales of Chrysophyceae in the recent history of Hall Lake, Washington. – *Freshwater Biology* 10: 61–66.
- NĚMCOVÁ, Y., NEUSTUPA, J., NOVÁKOVÁ, S. & KALINA, T. (2003): Silica-scaled chrysophytes of the Czech Republic. – *Act. Univ. Carol. Biol.* 47: 285–346.
- NOVÁKOVÁ, S., NĚMCOVÁ, Y., NEUSTUPA, J., ŘEZÁČOVÁ, M. & ŠEJNOHOVÁ, L. (2004): Silica-scaled chrysophytes in acid peat bogs of Bohemian Switzerland (Czech Republic) and Saxonian Switzerland (Germany). – *Nova Hedwigia* 78: 507–515.
- PICHRTOVÁ, M., ŘEZÁČOVÁ-ŠKALOUDOVÁ, M. & ŠKALOUD, P. (2007): The silica-scaled chrysophytes of the Czech-Moravian Highlands. – *Fottea, Olomouc*, 7: 43–48.
- PICHRTOVÁ, M. & VESELÁ, J. (in press): Silica-scaled chrysophytes in the Elbe Sandstone Region, Czech Republic. – *Fottea*.
- ŘEZÁČOVÁ, M. (2006): *Mallomonas kalinae* (Synurophyceae), a new species of alga from northern Bohemia, Czech Republic. – *Preslia* 78: 353–358.
- ŘEZÁČOVÁ, M., NEUSTUPA, J. & ŠEJNOHOVÁ, L. (2004): Five species of *Mallomonas* (Synurophyceae) new to the algal flora of the Czech Republic. – *Preslia* 76: 175–181.
- ŘEZÁČOVÁ, M. & NEUSTUPA, J. (2007): Distribution of the genus *Mallomonas* (Synurophyceae) – ubiquitous dispersal in microorganisms evaluated. – *Protist* 158: 29–37.
- SANDGREN, C.D. (1988): The ecology of chrysophyte flagellates: their growth and perennation strategies as freshwater phytoplankton. – In: SANDGREN, C.D. (ed.): Growth and reproductive strategies of freshwater phytoplankton. – pp. 9–104, Cambridge University Press, Cambridge.
- SIVER, P.A. (1991): The biology of *Mallomonas*. Morphology, taxonomy, ecology. – 230 pp., Kluwer Academic Publishers, Dordrecht.
- SIVER, P.A. (1995): The distribution of chrysophytes along environmental gradients: their use as biological indicators. – In: SANDGREN, C.D., SMOL, J.P. & KRISTIANSEN, J. (eds): Chrysophyte algae. – pp. 232–268, Cambridge University Press, Cambridge.
- SMOL, J.P. (2005): Applications of chrysophytes to problems in paleoecology. – In: SANDGREN, C.D., SMOL, J.P. & KRISTIANSEN, J. (eds): Chrysophyte algae. – pp. 232–268, Cambridge University Press, Cambridge.
- TAKAHASHI, E. (1959): Studies on genera *Mallomonas*, *Synura*, and other plankton in fresh-water by electron microscope (1). – *Bull. Yamagata Univ. Agr. Sci.* 3: 117–51.

© Czech Phycological Society

Received December 30, 2008

Accepted January 28, 2009