


University of South Bohemia

Faculty of Natural Sciences

Department of Molecular Biology and Biochemistry



Ph.D. Thesis

Aspects of RNA editing in *Trypanosoma brucei*

Mir Mohamod Hassan Hashimi

Supervisor: Prof. RNDR. Julius Lukeš

Institute of Parasitology, Biology Center, Czech Academy of Sciences

Laboratory of Molecular Biology of Protists



České Budějovice, Czech Republic, 2009

Hashimi, H., 2009: Aspects of RNA editing in *Trypanosoma brucei*. Ph.D. thesis, in English – 139p. Faculty of Natural Sciences, University of South Bohemia, České Budějovice.

Annotation:

This work addresses various aspects of RNA editing, a unique processing event that is essential for mitochondrial (mt) RNA maturation in the model kinetoplastid *Trypanosoma brucei*. RNAi-mediated reverse genetics was used extensively throughout this work. TbRGG1 appears to have a role in this process or stabilizing edited RNAs and is associated with a putative complex we called mt RNA binding complex 1. Several subunits have been demonstrated to have various roles in RNA metabolism, guide RNA biogenesis being the most significant. The extensively edited ATP subunit 6 was shown to be incorporated into the F_0F_1 ATP synthase complex by indirect genetic means. This respiratory complex is essential in both insect and infectious bloodstream stages of the parasite. Work here supports the idea that *Trypanosoma evansi* and *Trypanosoma equiperdum*, pathogens of livestock that are RNA-editing incompetent, should be considered petite mutants of *T. brucei*.


Financial support:


This work was supported by grants from the Grant Agency of the Czech Republic (204/06/1558 and 204/09/1667) (to J.L.) (524/03/H133) (to H.H.), Grant Agency of the Czech Academy of Sciences (A500960705) (to J.L.), National Institutes of Health (5R03TW6445) (K.D.S. and J.L.) (AI14102) (to K.S.) and Grant Agency of the Slovak Academy of Sciences (1/3241/06) (to A.H).

I hereby declare that I did all the work summarized in this thesis on my own or in collaboration with the co-authors of the presented papers and manuscripts, and only using the cited literature.

Acknowledgements

When I heard the word trypanosome, just as I was entering the Laboratory of Molecular

Biology of Protists in 2003, I pictured in my head  and was surprised to

see  wriggling throughout the microscope field of my first culture. It's probably not an understatement to write that I have learned a lot since that time, which I hope is somehow reflected in the thesis you have in your hands.

I would first like to thank the man who gave me this opportunity, Jula Lukeš, a true enthusiast whose energy has inspired a very stimulating and exciting environment. He is really one of those true personalities who enriched my life. Thank you for your support, Jula. I consider you to be a mentor and a friend.

I give a lot of heartfelt thanks to collaborators I have had the pleasure to work with over these years. I especially thank the members of a nice working group I (tried to) manage: Zdenka Čičová, Lucka Novotná, Yan-Zi Wen, Lucy Hanzálková and Jula Lukeš IV. Their efforts have greatly contributed to the manuscript in Chapter 3. Teaching you guys has been the most rewarding part of my PhD studies. I really thank you all for your patience with a guy struggling with the responsibilities of guiding a project and working group (a position that really made me appreciate how Jula keeps that full head of hair; I seem to be losing mine in a much lesser position), especially one whose priorities seem to change with each new meeting and publication.

Special thanks go to you Zdenka. We've been through a lot together. Another understatement? I've also learned a lot from you, and I hope that I somehow imparted some useful knowledge to you as well. I hope that fortune will shine on your future endeavors so that your success matches your potential.

It has been a privilege for me to spend 3 months in the laboratory of Ken Stuart at SBRI in Seattle. What an amazing group of people! I really learned so much from each member of that dynamic group, especially the PI, Achim Schnauffer, Reza Salavati, Nancy Ernst, Jason Carnes, James Trotter, Salvador Tarun, Rachel Dalley, Rose Proff,

Michele Fleck and Brian Panicucci, who gave me a sound foundation on how to conduct biological research. I thank each one and particularly Ken, Alena Zíková and Aswini Panigrahi, who were collaborators on the manuscript in Chapter 2.

Alena is a special case. I remember when we met at U Havrana and discussed what I needed to do to enter the University of South Bohemia over a few beers. She has constantly played the role of advisor to me, since she was a PhD student in Jula's lab to her current position in Ken's lab. Thanks Alena, and I look forward to bugging you for advice when you'll be next door in Alena's lab.

De-Hua Lai came to our lab in 2006 and was the driving force of the study of dk/ak cells resulting in manuscript in Chapter 3. Thanks for the opportunity to contribute to such an interesting topic.

I have the great pleasure of collaborating with Anton Horváth on what has been the never-ending manuscript. Aren't you happy I convinced you to work on this project? He and his studentky Petra Čermaková and Vladka Benkovičová lent their expertise to the manuscript in Chapter 5. Thanks for your belief in the project and stimulating discussions. I hope we can find another, more straightforward project to work on together.

I would like to thank past and present members of the Laboratory of Molecular Biology of Protists, especially Zdeňek Paris, Zdeňek Verner, Silva Foldynová-Trantírková, Evička Horáková, Daša Folková and Shao Long, for their help and support. I thank members of the Department of Molecular Biology for always being willing to lend valuable advice, in particular Ivoš Šauman, Lukaš Trantírek, Masako Asahina-Jindrová, Marek Jindra and Mira Oborník, as well as members of their respective labs. I also thank Tomáš Scholz and Libor Grubhoffer, director of the Institute of Parasitology and Dean of the Faculty of Natural Sciences, for their moral and financial support.

Great thanks go to the reviewers of this thesis, Achim, Laurie Read and Jan Tachezy. I am honored that you took the time from your busy schedules to do this. I look forward to your input.

Special thanks go to Barůška for her love, patience and support not just during my studies, but our life together. You've given me so much.

~hassan

Contents

Abbreviations

Summary

Chapter 1:

General introduction

Chapter 2:

TbRGG1, an essential protein involved in kinetoplastid RNA metabolism that is associated with a novel multiprotein complex (reprint of RNA *14*, 970-980).

Chapter 3:

Kinetoplastid guide RNA biogenesis is dependent on subunits of the mitochondrial RNA binding complex 1 and mitochondrial RNA polymerase (manuscript of RNA article *in press*).

Chapter 4:

Adaptations of *Trypanosoma brucei* to gradual loss of kinetoplast DNA: *Trypanosoma equiperdum* and *Trypanosoma evansi* are petite mutants of *T. brucei* (reprint of Proc. Natl. Acad. Sci. USA 105, 1999-2004).

Chapter 5:

The protein product of pan-edited ATP synthase subunit 6 is incorporated into a non-canonical Complex V in *Trypanosoma* Species (manuscript in preparation).

Conclusions and perspectives

Abbreviations

$\Delta\Psi_m$	mitochondrial membrane potential
A	adenosine
A6	F ₁ /F ₀ -ATP synthase subunit 6
ak	akinetoplastic
base "J"	beta-D-glucosyl-hydroxymethyluracil
CBP	calmodulin binding protein
complex I	NADH dehydrogenase
complex III	cytochrome <i>bc</i> ₁ complex
complex IV	cytochrome <i>c</i> oxidase
complex V	F ₁ /F ₀ -ATP synthase
coxX, coX	cytochrome <i>c</i> oxidase subunit X
cyB	cytochrome <i>b</i>
dk	dyskinetoplastic
dsRNA	double stranded RNA
G	guanosine
GAP	guide RNA associated protein
GAPDH	glyceraldehyde-3-phosphate
GRBC	guide RNA binding complex
gRNA	guide RNA
HA	hemagglutinin
HAT	human African trypanosomiasis
kDNA	kinetoplast DNA
KPAP	poly(A) polymerase
KREL	kinetoplastid RNA editing ligase
KREN	kinetoplastid RNA editing nuclease
KREP	kinetoplastid RNA editing proteins
KRET	kinetoplastid RNA editing 3' terminal uridylyltransferase
KREX	kinetoplastid RNA editing exonuclease
LSU	large subunit of riosome
mt	mitochondrial
MRB1	mitochondrial RNA binding complex 1

MRP	mitochondrial RNA binding protein
MURF	mitochondrial unidentified reading frame
NDx	NADH dehydrogenase subunit X
nt	nucleotide
Nudix	<u>n</u> ucleoside <u>d</u> iphosphate linked to some other moiety <u>X</u>
ORF	open reading frame
PARP	procyclic acidic repetitive protein
PPR	pentatricopeptide repeat
qPCR	quantitative real-time PCR
RNAi	RNA interference
RNAP	RNA polymerase
RPS12	ribosomal protein S12
SL RNA	spliced leader RNA
SSU	small subunit of ribosome
T	thymidine
TAO	trypanosome alternative oxidase
TAP	tandem affinity purification
tet	tetracycline
tetR	tetracycline repressor
U	uridine
UTR	untranslated region
VSG	variable surface glycoprotein

Summary

Members of the *Trypanosoma brucei* complex are the causative agents of several diseases, including human African trypanosomiasis and a wasting disease of ruminants called Nagana. This species has become a model species of its order Kinetoplastida, comprised of evolutionarily divergent flagellate protists, since it is amenable to RNA interference (RNAi). Several unique features of these cells include polycistronic nuclear transcription and *trans*-splicing, an extensive mitochondrial (mt) genome called kinetoplast (k) DNA and RNA editing. This thesis concerns aspects of RNA editing, an essential process in the maturation of transcripts encoded in kDNA maxicircles. It can be conceptually divided into two parts.

The first part (Chapters 2 and 3) deals with the functional analysis of proteins and protein complexes that are indirectly involved in RNA editing. RNAi-silencing of TbRGG1 showed that edited RNAs were downregulated. Steady state levels of guide (g) RNAs, vital genetic elements of this process encoded on kDNA minicircles, were not affected. Tandem affinity purification (TAP) of a TAP-tagged version of TbRGG1 over-expressed *in vivo* led to the discovery of a putative ~14 protein complex, provisionally named mitochondrial RNA binding complex 1 (MRB1). Three MRB1 subunits, named RNA helicase and gRNA associated proteins (GAPs) 1 and 2 have a role in some aspect of gRNA biogenesis or stability. The GAPs appear to be interacting partners, are essential in the bloodstream stage and have an interesting, punctuate localization in the mitochondrion of the procyclic stage. Downregulation of another MRB1 subunit, a predicted Nudix hydrolase, results in a general destability of maxicircle encoded RNAs without affecting gRNAs. A single mt RNA polymerase seems to transcribe both maxicircle and minicircle transcripts.

The second part (Chapters 4 and 5) deals with understanding the phenomena of dyskinetoplasty, in which the homogenization of minicircles leads to RNA editing incompetence and eventual loss of kDNA. *Trypanosoma evansi* and *Trypanosoma equiperdum*, important veterinarian pathogens, occur world-wide despite their aberrant kDNA. Analysis of these cells employing molecular and classical-parasitology approaches indicated that they are strains of *T. brucei* and should be given subspecies

status and considered petite mutants that are locked in the bloodstream stage. This state appears to arise spontaneously in perhaps a regular and/or autonomous fashion. However, their existence is not consistent with the finding that RNA editing is essential in the bloodstream stage of *T. brucei*, presumably for the extensive editing of a transcript provisionally assigned to be F₀F₁ ATP synthase subunit 6 (A6). This respiratory complex is essential in the bloodform to maintain membrane potential of the mitochondrion. A simple but unlikely explanation for dyskinetoplasty is that the mRNA in question does not encode A6 or that the protein product is not assembled into the complex. Resorting to an indirect, reverse genetics approach, since direct detection of almost all mt encoded proteins is not feasible, we provide evidence that A6 is indeed assembled into the F₀F₁ ATP synthase. Our results suggest that this complex may have an unusual structure as compared to those from other aerobic eukaryotes.

Chapter 1:

General Introduction – Table of Contents

Section	Page
1. The Kinetoplastida	1
2. The <i>Trypanosoma brucei</i> complex	2
3. A brief discussion of some unusual biological properties of <i>Trypanosoma brucei</i>	5
4. The extraordinary mitochondrion of <i>Trypanosoma brucei</i>	6
4.1 Atypical energy metabolism	6
4.2 Kinetoplast DNA	8
5. RNA editing in the mitochondrion of <i>Trypanosoma brucei</i>	10
5.1 Mechanism of RNA editing	10
5.2 The 20S editosome	12
5.3 Other proteins involved in RNA editing	15
5.4 The mitochondrial RNA complex 1	16
5.5 The <i>raison d'etre</i> of RNA editing	18
6. The mitochondrial RNA metabolism of <i>Trypanosoma brucei</i>	19
7. Translation of mitochondrial RNA	20
8. Dyskinetoplastic <i>Trypanosoma brucei</i>	21
9. Tools for functional analysis of <i>Trypanosoma brucei</i>	23
References	27

Figure	Page
1. Life cycle of <i>T. brucei</i> .	3
2. Physiological changes between bloodstream and procyclic stage mitochondria.	7
3. The complexity of the kDNA network.	9
4. Organization of <i>T. brucei</i> maxicircle genes.	10
5. Pan-editing of <i>cox3</i> .	11
6. Guide RNAs mediate RNA editing.	12
7. Mechanism of RNA editing.	13
8. Composition of 20S editosome.	14
9. The varying compositions of the putative mitochondrial RNA complex.	16
10. Structure of complex V.	23
11. Genetic background of 29-13 cell line.	24
12. pLew79-MHTAP cassette map.	26

Chapter 2:

TbRGG1, an essential protein involved in kinetoplastid RNA metabolism that is associated with a novel multiprotein complex. (reprint of Hashimi *et al.*, 2008, *RNA* 14, 970-980)

Abstract:

The uridine insertion / deletion RNA editing of kinetoplastid mitochondrial transcripts is performed by complex machinery involving a number of proteins and multiple protein complexes. Here we describe the effect of silencing of TbRGG1 gene by RNA interference on RNA editing in procyclic stage of *Trypanosoma brucei*. TbRGG1 is an essential protein for cell growth, the absence of which results in an overall decline of edited mRNAs, while the levels of never-edited RNAs remain unaltered. Repression of TbRGG1 expression has no effect on the 20S editosome and MRP1/2 complex. TAP-tag purification of TbRGG1 co-isolated a novel multi-protein complex and its association was further verified by TAP-tag analyses of two other components of the complex. TbRGG1 interaction with this complex appears to be mediated by RNA. Our results suggest that the TbRGG1 protein functions in stabilizing edited RNAs or editing efficiency and that the associated novel complex may have a role in mitochondrial RNA metabolism. We provisionally name it putative mitochondrial RNA binding complex 1 (put-MRB complex 1).

Chapter 3:

Kinetoplastid guide RNA biogenesis is dependent on subunits of the mitochondrial RNA binding complex 1 and mitochondrial RNA polymerase.

(manuscript of RNA article *in press*)

Kinetoplastid guide RNA biogenesis is dependent on subunits of the mitochondrial RNA binding complex 1 and mitochondrial RNA polymerase

Hassan Hashimi, Zdeňka Čičová, Lucie Novotná, Yan-Zi Wen[#] & Julius Lukeš^{*}

Biology Centre, Institute of Parasitology and Faculty of Natural Sciences, University of South Bohemia, České Budějovice (Budweis), Czech Republic

[#]Permanent address: Center for Parasitic Organisms, School of Life Sciences, Sun Yat-Sen (Zhongshan) University, Guangzhou 510275, P. R. China

^{*}Corresponding author: Institute of Parasitology, Branišovská 31, 37005 České Budějovice, Czech Republic; Tel: 00420-38-7775416; Fax: 0042-38-5310388; e-mail: jula@paru.cas.cz

Abstract

The mitochondrial RNA binding complex 1 (MRB1) is a recently discovered complex of proteins associated with the TbRGG1 and TbRGG2 in *Trypanosoma brucei*. Based on the phenotype caused by downregulation of these two proteins, it was proposed to play an unspecified role in RNA editing. RNAi-silencing of three newly characterized protein subunits, guide RNA associated proteins (GAPs) 1 and 2 as well as a predicted DExD/H-box RNA helicase, show they are essential for cell growth in the procyclic stage. Furthermore, their downregulation leads to inhibition of editing in only those mRNAs for which minicircle-encoded guide (g) RNAs are required. However, editing remains unaffected when the maxicircle-encoded *cis*-acting gRNA is employed. Interestingly, all three proteins are necessary for the expression of the minicircle-encoded gRNAs. Moreover, downregulation of a fourth assayed putative MRB1 subunit, Nudix hydrolase, does not appear to destabilize gRNAs and down-regulation of this protein has a general impact on the stability of maxicircle-encoded RNAs. GAP1 and 2 are also essential for the survival of the bloodstream stage, in which the gRNAs become eliminated upon depletion of either protein. Immunolocalization revealed that GAP1 and 2 are concentrated into discrete spots along the mitochondrion, usually localized in

proximity of the kinetoplast. Finally, we demonstrate that the same mt RNA polymerase known to transcribe the maxicircle mRNAs may also have a role in expression of the minicircle-encoded gRNAs.

Keywords: RNA editing, guide RNA, mitochondrion, trypanosome

Introduction

Kinetoplastid flagellates, the most studied of which are causative agents of African sleeping sickness, Chagas disease and leishmaniases in humans, have a strikingly complex mitochondrial (mt) DNA, also termed kinetoplast (k) DNA. It is composed of two types of molecules – maxicircles and minicircles. In the single mitochondrion, maxicircles are present in about a dozen of copies and represent the equivalents of classical mt DNA, bearing genes for subunits of respiratory complexes, mitoribosomal RNAs and one mitoribosomal subunit (for recent review see Simpson et al., 2004; Lukeš et al., 2005; Stuart et al., 2005). A unique feature of these flagellates is that the transcripts of some of the maxicircle-coded genes are post-transcriptionally altered by additions and/or deletions of uridylate (U) residues by a process termed RNA editing (Benne et al., 1986).

The function of kDNA minicircles had remained mysterious for decades. These molecules are concatenated in a network, number in the thousands, and their heterogenous sequence make up the bulk of the mt genome (Liu et al., 2005). The seminal discovery that they encode genes for small RNA molecules called guide (g) RNAs revealed their biological role as a genetic reservoir complementing the maxicircles (Sturm and Simpson, 1990). It is these gRNAs that provide information for multiple and specific insertions and deletions of uridines into pre-edited maxicircle mRNAs. Each gRNA is composed of three regions, differing in their function and content. The 5' anchor region hybridizes to a complementary sequence of the pre-edited mRNA, while the downstream information region of the RNA molecule provides the template for U insertions/deletions in the cognate mRNA via both Watson-Crick and non-canonical G:U base pairing. The 3' end of the gRNA is comprised of the post-transcriptionally added oligo(U) tail (Blum and Simpson, 1990). While the identity of the RNA components of editing was established almost 20 years ago, the identification of the myriad of proteins involved in mt RNA metabolism remains an ongoing endeavor.

The core RNA editing process has been ascribed to the 20S editosome, a multiprotein complex composed of 20 proteins (Simpson et al., 2004; Lukeš et al., 2005; Stuart et al., 2005). It has been firmly established by *in vitro* and *in vivo*

experiments that editing is carried out by this complex, which contains all catalytic activities necessary for the cleavage of the pre-mRNA, addition or deletion of the U(s), and ligation of the processed RNA molecules. The 20S editosome appears to be represented by a population of at least three dynamic protein complexes slightly differing in their composition and specific role in this process (Panigrahi et al., 2006; Carnes et al., 2008).

Mitochondrial RNA binding proteins have been a focus of intense research. The MRP1/2 complex is a heterotetrameric structure, composed of the MRP1 and MRP2 proteins, with positively charged residues concentrated at one face, to which the gRNAs anneal (Schumacher et al., 2006). Data obtained with the *in vivo* purified as well as *in vitro* reconstituted MRP1/2 complexes are consistent with a role in matchmaking the gRNAs with cognate pre-edited mRNAs (Muller et al., 2001; Zikova et al., 2008). RBP16 is another RNA binding protein that appears to have an *in vivo* role in editing (Pelletier and Read, 2003), perhaps via its demonstrated ability to stimulate insertion editing *in vitro* (Muller et al., 2006).

The initial characterization of TbRGG1, so-called because it possesses the RGG RNA binding motif, suggested the existence of another protein complex with a role in RNA editing (Vanhamme et al., 1998). The downregulation of TbRGG1 in the procyclic (=insect) stage of *T. brucei* caused an overall decline of edited mRNAs, but the never-edited transcripts were unaffected (Hashimi et al., 2008). This protein is stably associated with a putative protein complex composed of about 14 different subunits (Hashimi et al., 2008; Panigrahi et al., 2008), provisionally named the mitochondrial RNA binding complex 1 (MRB1 complex). Several of its components have features that point to RNA processing activities associated in an RNA mediated manner (Hashimi et al., 2008; Panigrahi et al., 2008). Another component of this complex, TbRGG2 (also known as TbRGGm), has affinity for poly(U) and its depletion leads to dramatic decrease of edited mRNAs (Fisk et al., 2008). Interestingly, a recent report on the newly discovered KPAP1 protein, poly(A) polymerase that polyadenylates maxicircle transcripts, shows that it is associated with a complex of proteins of overlapping composition with the MRB1 complex (Etheridge et al., 2008; Weng et al., 2008).

From these data, the role of the MRB1 complex is not obvious. However, its importance is underscored by the fact that orthologs of its subunits are found in all trypanosomatid genomes sequences thus far. Here, we attempt to ascertain the function of the MRB1 complex by functional analysis of four subunits of this complex by RNAi-silencing in the procyclic stage. The examined subunits are summarized in Table 1, including a nomenclature introduced in a similar study performed by Weng et al. (2008). Two of these subunits, Tb927.2.3800 and Tb927.7.2570, are paralogs that have no known protein motifs or domains (Hashimi et al., 2008). They will be referred to herein as guide RNA associated

proteins (GAP) 1 and 2, respectively, since we provide evidence for their participation in the biogenesis of these molecules. The third subunit, Tb927.4.1500, is predicted to be a ~240 kDa protein with DExD/H-box RNA helicase domains (Hashimi et al., 2008), and is referred to as such in this paper. The last studied subunit, Tb927.11.7290, is annotated as a Nudix hydrolase due to its conservation to other such proteins. Moreover, we show that mt RNA polymerase (mtRNAP), known to transcribe the maxicircle-encoded protein-coding genes (Grams et al., 2002), appears to have a similar role for minicircle-encoded gRNAs.

Materials and Methods

Generation of RNAi-knockdown and transgenic cell lines – Primers for generation of 400-600 bp gene fragments for cloning into the p2T7-177 vector were designed using the RNAit online tool available on the TrypanoFAN website (<http://trypanofan.path.cam.ac.uk/software/RNAit.html>). The following primer pairs (with restriction site underlined and indicated in parenthesis) were used: GAP1 gene – GAP1-F (5'-CTCCTCGAGCCTTTCAGC-3') (XhoI) and GAP1-R (5'-GGCAAGCTTCTGCGAATGTAG-3') (HindIII), amplified a 533 bp- long fragment; GAP2 gene – GAP2-F (5'-CGAGGATCCACAACGGCATT -3') (BamHI) and GAP2-R (5'-CATCATCAGCAAGCTTTATGATG-3') (HindIII), amplified a 580 bp-long fragment; He1-F (5'-GGATCCGTAGGAACTGGCAGAGACGC-3') (BamHI) and He1-R (5'-CTCGAGGCTACTGATTGCACGCAAAA -3') (XhoI), amplified a 453 bp-long fragment; Hyd1-F (5'-GGATCCCGTAAGGTGTCAGGACCGAT-3') (BamHI) and Hyd1-R (5'-CTCGAGTCTACGGTAATGCCCGTTTC-3') (XhoI) amplified a 467 bp-long fragment. All amplified fragments were cloned into the p2T7-177 vector creating constructs, which were upon linearization electroporated into the procyclic *T. brucei* strain 29-13, and the clones were obtained by limiting dilution at 27°C as described elsewhere (Vondrušková et al., 2005). Synthesis of dsRNA was induced by the addition of 1 µg/ml tetracycline to six clonal cell lines for each knockdown. For further experiments clones were selected based on the tightness of tetracycline expression of dsRNA and the robust elimination of target mRNA, as determined by Northern blot analysis using the appropriate gene fragment as a probe. The procyclic knockdown generated by cloning a fragment of the mt RNA polymerase gene in the pZJM vector (Grams et al., 2002) was kindly provided by P.T. Englund (Johns Hopkins University, Baltimore).

Linearized p2T7-177 vectors containing the GAP1 and 2 gene fragments were electroporated using the Amaxa Nucleofector II electroporator into the 427 bloodstream *T. brucei* strain. About 3×10^7 cells of exponentially growing culture were used for transfection, following the protocol of Vassella et al.(2001) with

minor modifications. Transfectants were selected with 1.25 µg/ml phleomycin and cloned by limiting dilution. RNAi was triggered and assessed in six clonal cell lines by the addition of 1 µg/ml tetracycline using the same strategy as that described for the procyclics.

The full-length open reading frames of the GAP1 and 2 genes were cloned into the pJH54 plasmid containing the C-terminal HA₃-tag (a gift from Christine Clayton, University of Heidelberg). The PCR amplified insert was digested with either HindIII and XbaI (GAP1) or just XbaI (GAP2). The orientation of the latter insert was verified by diagnostic restriction digests and sequencing. The utilized PCR primers are (with restriction site underlined): GAP1-HA-F (5'-AAGCTTATGCTGCGCGCGCCTG-3') and GAP1-HA-R (5'-TCTAGAGTATGCCGAAACGGCAGT-3'); GAP2-HA-F (5'-TCTAGAATGCTTCGCTTATTGCGG-3') and GAP2-HA-R (5'-TCTAGACAACTTCGCCTCACAGCC-3').

Generation of antibodies – Affinity purified polyclonal rabbit antibodies against GAPs 1 and 2 were raised against synthetic oligopeptides by the GenScript corporation (New Jersey, USA). The GAP1 synthetic oligopeptide (EYGDWGAEPGFEDRC) corresponds to the 137-150 amino acid region of the *T. brucei* protein while the GAP2 oligopeptide (TIKRSKDQLHVDLDC) is derived from the 297-310 amino acid region.

Western and Northern blot analyses – For Western blot analysis, cell lysates corresponding to 5 x 10⁶ procyclics/lane or 1 x 10⁷ bloodstreams/lane were separated on a 12% SDS-PAGE gel. The polyclonal rabbit antibodies against the MRP2 protein (Vondrušková et al., 2005), enolase (provided by P.A.M. Michels) and frataxin (Long et al., 2008) were used at 1:2000, 1:150,000 and 1:1000 dilutions, respectively. The antibodies against GAP1 and GAP2 were used at the respective dilutions 1:2000 and 1:500. Monoclonal antibodies against KREPA2, KREL1 and KREPA3 and Tbmp45 were used as described elsewhere (Panigrahi et al., 2001; Madison-Antenucci et al., 1998). Commercial antibodies against the HA₃ tag were used according to manufacturer's protocol (Sigma).

For Northern blot analysis, ~ 10 µg per lane of total RNA from procyclic or bloodstream cells was loaded on a 1% formaldehyde agarose gel, blotted and cross-linked as published elsewhere (Vondrušková et al., 2005). After pre-hybridization in NaPi solution (0.25M Na₂HPO₄ and 0.25M NaH₂PO₄, pH 7.2, 1 mM EDTA, 7% SDS) for 2 hrs at 55 °C, hybridization was performed overnight in the same solution at 55 °C. A wash in 2x SSC + 0.1% SDS at room temperature (RT) for 20 min was followed by two washes in 0.2x SSC + 0.1% SDS for 20 min each at 55 °C.

Growth curves – Following Northern blot analysis of several clones for each knockdown, one clone was selected for further experiments. Growth curves obtained over a period of 14 days in the presence or absence of the RNAi-induction agent tetracycline for transfected procyclics or 8 days for transfected bloodstreams. Cell density was measured every 24 hours using the Beckman Z2 Cell Counter.

Cell fractionation, glycerol gradients, digitonin fractionation – For separation on glycerol gradients, mitochondria from 10^9 procyclic cells were purified following the protocol detailed in Hashimi et al (2008). Digitonin fractionation and glycerol gradient sedimentation were performed as described elsewhere (Smid et al., 2006).

Quantitative real-time PCR and guanylyltransferase labeling – RNA for quantitative real-time (q) PCR and guanylyltransferase labeling were collected and DNased as described in Hashimi et al. (2008). After checking the integrity by running an aliquot on a formaldehyde gel, 4.5 μ g of RNA was used as a template for cDNA synthesis for subsequent qPCR analysis (see Hashimi et al., 2008 for details). A parallel cDNA synthesis reaction was set up with the same amount of RNA but without the reverse transcriptase to serve as a control for effective DNase treatment. Primers for maxicircle mRNAs, as well as the housekeeping β -tubulin and 18S rRNA transcripts serving as reference genes, are listed in Carnes et al. (2005) and Hashimi et al. (2008). New primer pairs amplifying cDNAs corresponding to transcripts of the subunits of the MRB1 complex are: GAP1-qPCR-Fw (5'-AACGTATTGCGGATGCTTAC-3') and Rv (5'-GAACCACACGCTCACAAACAG-3'); GAP2-qPCR-Fw (5'-CGGCTCATATTCCTGCCAATG-3') and Rv (5'-CGAAGTCCTCAGCAACCAACC-3'); Hel-qPCR-Fw (5'-ATCGCGTTAGGTGAAGCAGT-3') and Rv (5'-AAATGGGGATCCCTAAGGTG-3'); Hyd-qPCR-Fw (5'-ATTTTTCACCCTTGCACGTC-3') and Rv (5'-TCGATGGATTTGTTGTCACC-3'). Relative abundances of maxicircle transcripts were calculated using the Pfaffl method (Pfaffl, 2001).

Guanylyltransferase reactions were performed on 2.5-5 μ g of total RNA to cap gRNAs with [α - 32 P]GTP as previously described (Hashimi et al., 2008). The recombinant enzyme was either purchased from Ambion or provided by Ruslan Aphasizhev (University of California, Irvine). The reactions were separated on a denaturing 12% acrylamide-8M urea gel.

Immunolocalization – In order to visualize mitochondria, 5×10^6 to 1×10^7 cells were incubated in SDM-79 media supplemented with 1 mM of MitoTracker Green

FM (Molecular Probes) for 20 min at 27°C. Cells were subsequently washed 3x in phosphate-buffered saline (PBS) and then spread on slides, fixed and permeabilized as described in Klingbeil et al. (2002). All subsequent steps were carried out in a humid chamber. The slides were first blocked at RT for 1 hr with 10% goat serum diluted in PBS. Afterwards, the primary rabbit antibody was applied to the slides at the appropriate dilution in 10% goat serum + PBS for 90 min. After washing 3 x 10 min in PBS, the slides were incubated with secondary goat α -rabbit antibody conjugated to the cy3 fluorophore (Jackson ImmunoResearch) at a 1:800 dilution for 1 hr. The slides were then washed 3 x 10 min in PBS, and mounted in Vectashield containing DAPI. The α -GAP1 antibody was used at dilution 1:100 and the α -HA₃ antibody was used according to the manufacturer's protocol (Sigma). Samples probed with the latter antibody were not pretreated with MitoTracker. Cells were examined on the Olympus Fluoview FV1000 confocal microscope using the accompanying Fluoview v1.7 software. A composite of the Z-stack of the fluorescent images was rendered using the ImageJ software and processed using Adobe Photoshop v6.0.

Results

Subunits of the MRB1 complex are essential for the procyclic stage – Comparison of the growth of non-induced and RNAi induced GAP1, GAP2, RNA helicase and Nudix hydrolase cell lines revealed that these proteins are essential for the growth of *T. brucei* procyclics. In these cell lines, growth inhibition is apparent 3 to 4 days after addition of the RNAi-induction agent tetracycline (Fig. 1). In the case of the GAP knockdowns, the cells became increasingly resistant to the dsRNA and recovered around day 12 (Figs. 1A and 1B), as previously reported in the *T. brucei* system (Pelletier and Read, 2003). However, the RNA helicase and Nudix hydrolase knockdowns did not recover their wild-type growth over the 14-day time-course. Based on these data, the timepoints after 3 to 4 days of RNAi induction were selected for all subsequent experiments with the knockdowns.

Stability of GAPs1 and 2 are mutually dependent – To confirm the mutual association of the GAP proteins as predicted by the mass spectrometry analysis (Hashimi et al., 2008), the mt lysates of cells with a single down-regulated GAP were probed with polyclonal antibodies against both proteins. The specificity of each α -GAP antibody for the appropriate target protein was verified by probing recombinant GAPs overexpressed in *E. coli* (data not shown). Indeed, elimination of the target protein by RNAi was followed by the disappearance of the other GAP protein (Fig. 2A), testifying to their mutual dependence. Furthermore, we checked

whether the GAPs are also destabilized upon silencing of either the RNA helicase or Nudix hydrolase. In either of these backgrounds, both proteins persisted (Fig. 2B).

RNAi-silencing of MRB1 complex subunits affects maxicircle transcripts – Since TbRGG1 and TbRGG2 from the MRB1 complex were shown to be essential for RNA editing (Hashimi et al., 2008; Fisk et al., 2008), we decided to investigate in detail the effect of down-regulation of the studied subunits in procyclic cells on the stability and editing of mt-encoded mRNAs. The levels of several pre-edited, edited and never-edited RNAs were measured by a quantitative real time (q) PCR based assay, using cDNA from RNAi-induced and non-induced cells. Primers amplifying the pre-edited and edited mt mRNAs of ATPase subunit 6 (A6), cytochrome oxidase subunits 2 and 3 (cox2 and cox3), cytochrome reductase subunit B (cyB), maxicircle unknown reading frame 2 (MURF2), NADH dehydrogenase subunit 7 (ND7) and mitoribosomal protein S12 (RPS12) were used. The levels of two never-edited mRNAs (ND4 and cox1) and, in some cases, the 9S and 12S mitoribosomal RNAs were also measured, as were the mRNAs targeted by RNAi. The obtained values were normalized to the measured levels of the cytoplasmic β -tubulin and 18S rRNA transcripts, since they are unaffected by RNAi. All qPCR reactions with a given primer pair were done in triplicate, and the average and median standard deviations of the measured cycle threshold values are given in Fig. 3.

The qPCR assay revealed a virtually identical phenotype for both GAP1 and 2 knockdowns (Figs. 3A and 3B). With a single but important exception (see below), all examined pre-edited mRNAs (A6, cox3, cyB, MURF2, ND7 and RPS12) were up-regulated between 1.5 and up to 20-fold, as compared to RNA from the non-induced cells. Importantly, this up-regulation was correlated to the decrease of the respective edited transcripts. No effect was observed on the never-edited mRNAs (cox1 and ND4) and rRNAs (9S and 12S) (Figs. 3A and 3B). Significantly, pre-edited and edited cox2 mRNAs were also not affected by RNAi-mediated elimination of either GAP. This transcript is unique in that its single gRNA is contained in its 3'-untranslated region, acting as a guide for insertion editing in *cis* (Golden and Hajduk, 2005). The assay also reveals that dsRNAs specifically target only the intended GAP RNA (Figs. 3A and 3B).

This assay revealed a similar effect on maxicircle transcripts in cells with silenced RNA helicase: the combined increase and decrease of pre-edited and edited transcripts, respectively, with the exception of cox2 transcripts (Fig. 3C). However, it should be noted that while never-edited transcripts were unaffected, so were the levels of pre-edited and edited ND7. In contrast, silencing of Nudix hydrolase showed a general effect on almost all maxicircle-encoded transcripts (Fig. 3D). Most never-edited, pre-edited and edited mRNAs and rRNAs are

downregulated upon depletion of this protein. This phenotype represents a highly significant departure from the otherwise relatively uniform phenotype of the other examined knockdowns.

The MRB1 complex has a role in gRNA expression– The disruption of RNA editing, observed in cells in which either GAP1, GAP2 or RNA helicase are downregulated, appears to be independent of the core editing activities of the 20S editosome since *cox2* mRNA editing proceeds normally. Another component of this process are the small 30-60 nt gRNAs. To see whether these primary transcripts are affected, they were capped with guanylyltransferase and [α - 32 P]GTP and visualized on a high-resolution denaturing acrylamide gel. An upper band corresponding to the cytoplasmic 5S rRNA is used as a loading control. Indeed, repression of GAP1 and 2 as well as RNA helicase in the procyclic cells leads to a decrease of the steady-state level of gRNAs (Figs. 4A, 4B and 4C). In contrast, neither the abundance nor 3'-oligo(U) tail of gRNAs is not affected upon depletion of Nudix hydrolase (Fig. 4D).

GAP1 assembly into macromolecular complexes is affected in the knockdowns for subunits of the MRB1 complex– To further study the sedimentation properties of GAP1 in cells with down-regulated MRB1 subunits, lysates from hypotonically isolated mitochondria from non-induced and RNAi-induced cultures were loaded onto 10 to 30% glycerol gradients. Gradient fractions were first probed with monoclonal antibodies against three of the core editosome proteins, KREPA2, REL1 and KREPA3 (Panigrahi et al., 2001), used as markers for 20S. As an additional control, the fractions were immunodecorated with an antibody binding Tbmp45, a protein previously labeled REAP1, as a marker for 40S sedimentation (Madison-Antenucci et al., 1998; Fisk et al., 2008). A representative immunoblot performed on GAP1 induced and non-induced samples is in agreement with previous studies, as all proteins recognized by these antibodies localized to fractions 13 thru 17, the peak of 20S editosome sedimentation (Fig. 5A, top panel). Furthermore, it has been demonstrated that this complex is not disrupted when substrate RNAs are absent since it is held together by protein-protein interactions (Domingo et al., 2003; Fisk et al., 2008; Hashimi et al., 2008). The Tbmp45 protein was found to be localized in the lowest fractions, as expected (Fig. 5A, bottom panel). The other gradients depicted in Fig. 5B were also validated in this fashion (data not shown).

In the separated mt lysates from the non-induced cell lines, GAP1 is distributed throughout the gradient, although with some variation in the pattern as revealed by Western analysis (Figs. 5B and 7B). A broad distribution of GAP2 has been previously reported (Panigrahi et al., 2008). As expected, GAP1 is reduced

upon interference of expression of either GAP subunits (Fig. 5B). A dramatic shift of the S value of GAP1 is observed in the RNA helicase-silenced cells, in which it is concentrated in the lighter fractions (Fig. 5B). Such a striking difference in GAP1 sedimentation is not observed in the induced and non-induced Nudix hydrolase knockdowns (Fig. 5B). In the hydrolase-silenced sample, a reduction of GAP1 immunopositive signal in the dense fractions is apparent (Fig. 5B).

mtRNA polymerase appears to transcribe minicircles – The participation of a multiprotein complex in the transcription of minicircle gRNA genes, independent of mtRNAP, has been hypothesized (Grams et al., 2002). In the inducible mtRNAP knockdowns (kindly provided by P.T. Englund), we have first verified the RNAi effect by Northern analysis, as antibodies against the *T. brucei* mtRNA polymerase are not available. As shown in Fig. 6A, the mtRNA polymerase-encoding mRNA is undetectable 3 days after tetracycline induction, the time-point in which growth inhibition is first observed (Grams et al., 2002). Next, we have hybridized blots containing mt RNA with a probe against *cox1* mRNA. In agreement with the previous study (Grams et al., 2002), this never-edited transcript is undetectable 3 days after RNAi is triggered (Fig. 6B). To test whether this polymerase is responsible for the transcription of minicircles, the initial step in expression of gRNA molecules, we have performed the guanylyltransferase labeling assay on total RNA isolated from non-induced and induced knockdowns for mtRNA polymerase. Clearly, the gRNA population is decreased in cells lacking the enzyme (Fig. 6C).

MRB1 complex is disrupted upon downregulation of mtRNA polymerase– The similarity in the effect on gRNA expression of the knockdowns for mtRNAP and MRB1 subunits GAP1, GAP2 and RNA helicase inspired us to investigate the integrity of this complex in the absence of the polymerase. For this purpose, total lysates prepared from non-induced cells as well as from cells in which mtRNAP was targeted by RNAi for 3 days were loaded on glycerol gradients and the obtained fractions were screened with the α -GAP1 antibody. The quality of the gradient was tested by Western blot analysis using antibodies against the editosome KREL1 and *Tbmp45* (Fig. 7A). GAP1 is broadly distributed in the non-induced cells in fractions 7 to 17 (Fig. 7B). Indeed, an apparent shift towards the lighter density fractions (with a marked peak in fractions 7 and 9) of the gradient in the GAP1 distribution occurs when mtRNAP is downregulated. This result indicates that in the absence of mtRNAP, the MRB1 complex is disrupted in the same manner as in the absence of RNA helicase (Fig. 5B).

GAP1 and 2 and gRNAs are essential for the bloodstream stage – Since RNA editing is essential also in the bloodstream stage (Schnauffer et al., 2001), we decided to test whether the absence of GAP1 and 2, and consequently gRNA expression, is also lethal in this stage. The constructs containing the same GAP1 or 2 gene fragments as those used for RNAi in the procyclics were transfected into the bloodstream stage. Western blot analysis revealed that in the GAP1 and 2 knockdowns, the targeted protein was undetectable after 2 days of induction (Figs. 8A). Moreover, as in the procyclic stage, the ablation of GAP1 leads to the reduction of GAP2, and vice versa, testifying to their mutual dependence.

Downregulation of either GAP1 or 2 results in growth inhibition, indicating that these proteins are also essential in this life cycle stage. When compared with the non-induced cells, the growth of the induced clones decreases at day 3 and stops at day 4 (Fig. 8B). Both induced cell lines recovered from the effect of RNAi at day 7 post-induction. Labeling of total RNA collected at days 2 and 3 post-induction with guanylyltransferase revealed that steady state levels of gRNAs are decreased when either GAP1 or 2 are silenced in the bloodstream stage (Fig. 8C). This result indicates that the function of these proteins is conserved between the two major life stages of *T. brucei*, and that gRNAs are also required for survival of the infective bloodforms.

GAP1 and GAP2 have a novel localization within the mitochondrion – Previous data provided evidence for localization of the MRB1 complex in the mitochondrion (Hashimi et al., 2008; Panigrahi et al., 2008). We have therefore resorted to indirect immunofluorescence experiments to determine where these proteins are located in the organelle. Using the α -GAP1 antibody, we show that in the procyclic cells, the protein is confined to several discrete foci, distributed throughout the reticulated mitochondrion (Fig. 9A, top row), often in the proximity of the kDNA disk. Specificity of the used antibody was confirmed by its failure to detect target protein in RNAi-silenced GAP1 cells (Fig. 9A, bottom row). The polyclonal α -GAP2 antibodies proved to be less useful in the immunolocalization studies.

To corroborate this unexpected observation, cells were transfected with a construct for tetracycline-inducible expression of either GAP1 or 2 proteins bearing an HA₃-epitope tag. Immunocytochemistry and DAPI staining performed on cells grown in the presence of the antibiotic for 2 days confirmed that both epitope-tagged proteins are confined to multiple discrete punctuate loci unevenly distributed throughout the organelle (Fig. 9B). As observed with the α -GAP1 antibody, an immunopositive signal is also often observed near the kinetoplast.

Although attachment of the HA₃ tag to the C-terminus of the protein should not interfere with the mt import signal located at the N-terminus, possible extra-mitochondrial localization had to be ruled out. Therefore, the 29-13 procyclics

containing the HA₃-tagged GAP1 or 2 were fractionated by extraction with increasing concentrations of digitonin. Indeed, probing the obtained cytosolic and mt fractions, as well as the total cell lysate, with antibodies against the HA₃ tag showed that both tagged GAP proteins are properly imported into the organelle (Fig. 9C).

Discussion

Since the discovery of RNA editing in 1986 (Benne et al., 1986), considerable efforts have been directed at understanding the molecular mechanisms underlying what was initially an enigmatic biological process. A major breakthrough was the discovery of gRNAs, which revealed that the information for U insertion/deletion events reside in these molecules and a function of the thousands of minicircles is to encode them (Blum and Simpson, 1990). The establishment of an *in vitro* editing assay led to the validation that proteins provided the enzymatic machinery for RNA editing (Seiwert et al., 1996). The isolation and characterization of the 20S editosome have dissected how a multi-protein complex can orchestrate the catalytic steps required for the maturation of most mt RNAs in trypanosomes (Simpson et al., 2004; Lukeš et al., 2005; Stuart et al., 2005).

Studies focusing on identifying proteins involved in other aspects of mt RNA metabolism have uncovered the MRB1 complex in three different instances (Etheridge et al., 2008, Hashimi et al., 2008, Panigrahi et al., 2008). Although the subunit composition in these studies is not identical, the degree of overlap is quite significant. The MRB1 complex contains up to 14 proteins, some of which possess known motifs and domains involved in RNA processing or protein interactions (Hashimi et al., 2008; Panigrahi et al., 2008). Two components of the MRB1 complex have been already characterized to some extent, namely TbRGG1 and TbRGG2, the latter of which was suggested to be a transiently associated editing factor required for the processivity of RNA editing (Fisk et al., 2008). RNAi-silencing of both proteins leads to a downregulation of edited transcripts, suggesting some kind of a role in RNA editing (Fisk et al., 2008; Hashimi et al., 2008). However, these data do not adequately address a possible role of the MRB1 complex.

Here we addressed the question of function by examining the phenotype resulting from RNAi-silencing of four subunits of this complex: GAP1, GAP2, RNA helicase and Nudix hydrolase. We have shown that a subset of the MRB1 complex, GAP1, GAP2 and RNA helicase, is involved in gRNA biogenesis. This notion is supported by the absence of gRNAs when any of these proteins is depleted as a consequence of RNAi. Furthermore, editing of the maxicircle-

encoded RNAs is subsequently reduced, with the notable exception of *cox2*. This transcript contains its own *cis*-gRNA in the 3'-untranslated region, thus circumventing the requirement for minicircle-encoded gRNAs acting in *trans* (Golden and Hajduk, 2005). The independence of *cox2* mRNA editing, in the *trans*-acting gRNA-deficient background, on these MRB1 proteins clearly indicates that they are not involved in core RNA editing activities.

The question is specifically what role do these proteins play in the expression of gRNAs? It was hypothesized that a multiprotein complex may be involved in the transcription of minicircle-encoded genes, independent of the canonical mt RNA polymerase (mtRNAP) involved in maxicircle transcription (Grams et al., 2002). Here we provide evidence that mtRNAP is the most likely candidate for the transcription of gRNA genes as well as maxicircle genes. This observation indicates that the MRB1 complex may have a distinct role in gRNA synthesis, since mtRNAP was not demonstrated to be associated with this complex in any of the published reports (Etheridge et al., 2008, Hashimi et al., 2008, Panigrahi et al., 2008). Furthermore, the possibility that the mtRNAP transcribes both minicircle-encoded gRNAs and maxicircle-encoded mRNAs is surprising given the requirement of the multiple DNA ligases and DNA polymerases for kDNA maintenance and replication (Klingbeil et al., 2002; Liu et al., 2005).

The GAPs and RNA helicase do appear to be involved in some aspect of gRNA processing and/or stability. The most established processing event in the biogenesis of a gRNA is its post-transcriptional addition of the 3'-oligo(U) tail (Blum et al., 1991) by the KRET1 enzyme (Aphasizhev et al., 2003b). It was also proposed that functional gRNAs are spliced from the 5'-ends of polycistronic minicircle transcripts (Grams et al., 2000). However, a corresponding accumulation of these large precursor RNAs was not detected in the GAP knockdowns (data not shown), suggesting that if this processing step does exist, it is not mediated by these proteins.

Another proposed feature of gRNA is the presence of two intramolecular hairpin structures with single stranded ends (Schmid et al., 1995; Hermann et al., 1997). The presence of a predicted DExD/H-box RNA helicase in the gRNA-expression component of the MRB1 complex raises the possibility that these proteins may play a role in facilitating the formation of this secondary structure in these small transcripts. Perhaps these thermodynamically unfavorable loops (Schmid et al., 1995) are not formed in the absence of the GAPs or RNA helicase, leading to the apparent destabilization of the gRNAs. However, this speculative model is tempered by the involvement of the DExD/H box RNA helicases in diverse roles in RNA metabolism not restricted just to processing, such as transcription regulation and RNA turnover (reviewed by Fuller-Pace, 2006).

We show that the GAP proteins are essential for the bloodstream stage, where they appear to have the same role in gRNA processing/stability as they do in the procyclic stage. Although an active RNA metabolism has been established as a requirement for survival of this infective stage (Schnauffer et al., 2001; Fisk et al., 2008), to our knowledge this result is the first to indicate that the gRNA biogenesis is required in this part of the life cycle as well. This result is significant in light of the existence of naturally occurring dyskinetoplastic and akinetoplastic trypanosomes (Lai et al., 2008), which must circumvent the lack of the requisite repertoire of gRNAs for the expression of normally essential mt-encoded genes, such as the ATPase subunit 6 (Schnauffer et al., 2005). Apparently, in both stages all kDNA is transcribed, and any regulation occurs at the post-transcriptional level, a situation strikingly similar to what is known for the expression of nuclear genes in trypanosomes and other kinetoplastid flagellates (Haile and Papadopoulou, 2007).

We also establish here that the group of proteins that we designate to the MRB1 complex do not share an overall function. While GAP1, GAP2 and RNA helicase are involved in gRNA expression, TbRGG1 and 2 are clearly not (Hashimi et al., 2008; Fisk et al., 2008). Furthermore, we show here that the Nudix hydrolase is also not involved in gRNA expression, since these molecules persist in a background deficient of this protein. The family of Nudix hydrolases has a diverse array of functions in the cell, such as decapping mRNAs (Cohen et al., 2005) and clearance of oxidized nucleosides (reviewed by McClennan, 2006). Designation of a specific role of this hydrolase is still not feasible based on the observed destabilization of most of the maxicircle transcripts upon its depletion. However, these results counter those of Weng and colleagues (2008) indicating a specific decrease in edited RNAs.

The array of phenotypes affecting different aspects of mt RNA metabolism upon downregulation of various putative subunits of the MRB1 complex, along with the assortment of associating proteins, is quite striking (Etheridge et al., 2008; Fisk et al., 2008; Hashimi et al., 2008; Panigrahi et al., 2008; Weng et al., 2008). These observations raise the question whether it represents a single *bona fide* complex, held together by protein-protein interactions, or a collection of monomers and/or smaller complexes connected by substrate RNAs. Indeed, the latter situation has been proposed (Weng et al., 2008) and incorporation of both TbRGG proteins into large complex(es) appears to be RNase sensitive, while the 20S editosome complex remains intact (Fisk et al., 2008; Hashimi et al., 2008). Interestingly, RNAi-silencing of either RNA helicase or mtRNAP clearly disrupts the GAP1 assembly into macromolecular complexes. This result may reflect a genuine interaction of these proteins with GAP1 or be a result of the consequent decrease of RNA molecules via which these proteins seem to associate.

We conclude by discussing the unique localization of the GAP proteins to discrete points along the reticulated mitochondrion, often proximal to the kDNA network. To our knowledge, such a pattern has not been reported for other proteins involved in mt nucleic acid metabolism (Klingbeil et al., 2002; Liu et al., 2005; Vanhamme et al., 1998; Etheridge et al., 2008). Panigrahi and co-workers (2008) have previously reported a more uniform distribution of different subunits of the MRB1 complex throughout the mitochondrion. However, given the aforementioned diversity of RNAi-silencing phenotypes for the MRB1 complex proteins, it is possible that they may also exhibit diverse yet overlapping sites in the organelle as well. Nevertheless, as the GAP proteins have a demonstrated role in gRNA biogenesis, we imagine these loci may represent centers, where gRNAs are processed and distributed for their participation in RNA editing as catalyzed by the 20S editosome.

Acknowledgements

We thank Lucie Hanzálková and Julius Lukeš IV for help with some experiments, and Paul T. Englund (Johns Hopkins University, USA), Paul A.M. Michels (Université catholique de Louvain, Belgium), Ken Stuart (Seattle Biomedical Research Institute, USA), Stephen Hajduk (University of Georgia, USA), Christine Clayton (Heidelberg University, Germany) and Ruslan Aphasizhev (University of California, Irvine, USA) for sharing the mt RNA polymerase knockdowns, the α -enolase antibody, antibodies against 20S editosome subunits, α -Tbmp45 antibody, pJH54 plasmid and recombinant guanylyltransferase, respectively. We also thank Aswini Panigrahi and Alena Zíková (Seattle Biomedical Research Institute, USA) for critical reading of the manuscript. This work was supported by the Grant Agency of the Czech Republic 204/09/1667, Grant Agency of the Czech Academy of Sciences A500960705, and the Ministry of Education of the Czech Republic (LC07032 and 2B06129) to J.L. and the PhD student grant 524/03/H133 to H.H.

References

- Aphasizhev, R., Aphasizheva, I., Nelson, R.E., and Simpson, L. 2003a. A 100-kDa complex of two RNA-binding proteins from mitochondria of *Leishmania tarentolae* catalyzes RNA annealing and interacts with several RNA editing components. *RNA* **9**: 62-76.
- Aphasizhev, R., Aphasizheva, I., and Simpson, L. 2003b. A tale of two TUTases. *Proc. Natl. Acad. Sci. USA* **100**: 10617-10622.

Benne, R., van den Burg, J., Brakenhoff, J.P., Sloof, P., Van Boom, J.H., and Tromp, M.C. 1986. Major transcript of the frameshifted coxII gene from trypanosome mitochondria contains four nucleotides that are not encoded in the DNA. *Cell* **46**: 819-826.

Blum, B., Bakalara, N., and Simpson, L. 1990. A model for RNA editing in kinetoplastid mitochondria: "guide" RNA molecules transcribed from maxicircle DNA provide the edited information. *Cell* **60**: 189-198.

Blum, B., and Simpson, L. 1990. Guide RNAs in kinetoplastid mitochondria have a nonencoded 3' oligo(U) tail involved in recognition of the preedited region. *Cell* **62**: 391-397.

Carnes, J., Trotter, J.R., Ernst, N.L., Steinberg, A.G., and Stuart, K. 2005. An essential RNase III insertion editing endonuclease in *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. USA* **102**: 16614-16619.

Carnes, J., Trotter, J.R., Peltan, A., Fleck, M., and Stuart, K. 2008. RNA editing in *Trypanosoma brucei* requires three different editosomes. *Mol. Cell. Biol.* **28**: 122-130.

Cohen, L.S., Mikhli, C., Jiao, X., Kiledjian, M., Kunkel, G., and Davis, R.E. 2005. Dcp2 decaps m^{2,2,7} GpppN-capped RNAs, and its activity is sequence and context dependent. *Mol. Cell. Biol.* **25**: 8779-8791.

Domingo, G.J., Palazzo, S.S., Wang, B., Panicucci, B., Salavati, R., and Stuart, K.D. 2003. Dyskinetoplastic *Trypanosoma brucei* contains functional editing complexes. *Eukaryot. Cell* **2**: 569-577.

Etheridge, R.D., Aphasizheva, I., Gershon, P.D., and Aphasizhev, R. 2008. 3' adenylation determines mRNA abundance and monitors completion of RNA editing in *T. brucei* mitochondria. *EMBO J.* **27**: 1596-1608.

Fisk, J.C., Ammerman, M.L., Presnyak, V., and Read, L.K. 2008. TbRGG2, an essential RNA editing accessory factor in two *Trypanosoma brucei* life cycle stages. *J. Biol. Chem.* **283**: 23016-23025.

Fuller-Pace, F.V. 2006. DExD/H box RNA helicases: multifunctional proteins with important roles in transcriptional regulation. *Nucl. Acids Res.* **34**: 4206-4215.

Golden, D.E., and Hajduk, S.L. 2005. The 3'-untranslated region of cytochrome oxidase II mRNA functions in RNA editing of African trypanosomes exclusively as a cis guide RNA. *RNA* **11**: 29-37.

Grams, J., McManus, M.T., and Hajduk, S.L. 2000. Processing of polycistronic guide RNAs is associated with RNA editing complexes in *Trypanosoma brucei*. *EMBO J.* **19**: 5525-5532.

Grams, J., Morris, J.C., Drew, M.E., Wang, Z.F., Englund, P.T., and Hajduk, S.L. 2002. A trypanosome mitochondrial RNA polymerase is required for transcription and replication. *J. Biol. Chem.* **277**: 16952-16959.

Haile, S., and Papadopoulou, B. 2007. Developmental regulation of gene expression in trypanosomatid parasitic protozoa. *Curr. Opin. Microbiol.* **10**: 569-577.

Hashimi, H., Zíková, A., Panigrahi, A.K., Stuart, K.D., and Lukeš, J. 2008. TbRGG1, a component of a novel multi-protein complex involved in kinetoplastid RNA editing. *RNA* **14**: 970-980.

Hermann, T., Schmid, B., Heumann, H., and Göringer, H.U. 1997. A three-dimensional working model for a guide RNA from *Trypanosoma brucei*. *Nucl. Acids Res.* **25**: 2311-2318.

Klingbeil, M.M., Motyka, S.A., and Englund, P.T. 2002. Multiple mitochondrial DNA polymerases in *Trypanosoma brucei*. *Mol. Cell* **10**: 175-186.

Lai, D.-H., Hashimi, H., Lun, Z.-R., Ayala, F.J., and Lukeš, J. 2008. Adaptations of *Trypanosoma brucei* to gradual loss of kinetoplast DNA: *Trypanosoma equiperdum* and *Trypanosoma evansi* are petite mutants of *T. brucei*. *Proc. Natl. Acad. Sci. USA* **105**: 1999-2004

Liu, B.Y., Liu, Y.N., Motyka, S.A., Agbo, E.E.C., and Englund, P.T. 2005. Fellowship of the rings: the replication of kinetoplast DNA. *Trends Parasitol.* **21**: 363-369.

Long, S., Jirků, M., Mach, J., Ginger, M.L., Sutak, R., Richardson, D., Tachezy, J., and Lukeš, J. 2008. Ancestral roles of eukaryotic frataxin: mitochondrial frataxin function and heterologous expression of hydrogenosomal *Trichomonas* homologues in trypanosomes. *Mol. Microbiol.* **69**: 94-109.

Lukeš, J., Hashimi, H., and Zíková, A. 2005. Unexplained complexity of the mitochondrial genome and transcriptome in kinetoplastid flagellates. *Curr. Genet.* **48**: 277-299.

Madison-Antenucci, S., Sabatini, R.S., Pollard, V.W. and Hajduk, S.L. 1998. Kinetoplastid RNA-editing-associated protein 1 (REAP-1): a novel editing complex protein with repetitive domains. *EMBO J.* **21**: 6368-76.

McClennan, A.G. 2006. The Nudix hydrolase superfamily. *Cell. Mol. Life Sci.* **63**: 123-143.

Miller, M.M., Halbig, K., Cruz-Reyes, J. and Read, L.K. 2006. RBP16 stimulates trypanosome RNA editing *in vitro* at an early step in the editing reaction. *RNA* **12**: 1292-1303.

Müller, U. F., Lambert, L., and Göringer, H.U. 2001. Annealing of RNA editing substrates facilitated by guide RNA-binding protein gBP21. *EMBO J.* **20**: 1394-1404.

Panigrahi, A.K., Schnauffer, A., Carmean, N., Igo, R.P.Jr., Gygi, S.P., Ernst, N.L., Palazzo, S.S., Weston, D.S., Aebersold, R., Salavati, R., and Stuart, K. D. 2001. Four related proteins of the *Trypanosoma brucei* RNA editing complex. *Mol. Cell. Biol.* **21**: 6833-6840.

Panigrahi, A.K., Ernst, N.L., Domingo, G.J., Fleck, M., Salavati, R., and Start, K.D. 2006. Compositionally and functionally distinct editsomes in *Trypanosoma brucei*. *RNA* **12**: 1038-1049.

Panigrahi, A.K., Zíková, A., Halley, R.A., Acestor, N., Ogata, Y., Myler, P.J., and Stuart, K. 2008. Mitochondrial complexes in *Trypanosoma brucei*: a novel complex and a unique oxidoreductase complex. *Mol. Cell. Proteomics* **7**: 534-545.

Parish, S., Reich, W., and Moss, B. 2007. Vaccinia virus D10 protein has mRNA decapping activity, providing a mechanism for control of host and viral gene expression. *Proc. Natl. Acad. Sci. USA* **104**: 2139-2144.

Pelletier, M., and Read, L.K. 2003. RBP16 is a multifunctional gene regulatory protein involved in editing and stabilization of specific mitochondrial mRNAs in *Trypanosoma brucei*. *RNA* **9**: 457-468.

- Pfaffl, M. W. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucl. Acids Res.* **29**, e45.
- Schmid, B., Riley, G.R., Stuart, K. and Göringer, H.U. 1995. The secondary structure of guide RNA molecules from *Trypanosoma brucei*. *Nucl. Acids Res.* **23**: 3093-3102.
- Schnauffer, A., Panigrahi, A.K., Panicucci, B., Igo, R.P.Jr., Wirtz, E., Salavati, R., and Stuart, K. 2001. An RNA ligase essential for RNA editing and survival of the bloodstream form of *Trypanosoma brucei*. *Science* **291**: 2159-2162.
- Schnauffer, A., Clark-Walker, G.D., Steinberg, A.G., and Stuart, K. 2005. The F₁-ATP synthase complex in bloodstream stage trypanosomes has an unusual and essential function. *EMBO J.* **24**: 4029-4040.
- Schumacher, M.A., Karamooz, E., Zíková, A., Trantírek, L., and Lukeš, J. 2006. Crystal structures of *T. brucei* MRP1/MRP2 guide-RNA-binding complex reveals RNA matchmaking mechanism. *Cell* **126**: 701-711
- Seiwert, S.D., Heidmann, S., and Stuart, K. 1996. Direct visualization of uridylate deletion in vitro suggests a mechanism for kinetoplast RNA editing. *Cell* **84**: 831-841.
- Simpson, L., Aphasizhev, R., Gao, G., and Kang, X. 2004. Mitochondrial proteins and complexes in *Leishmania* and *Trypanosoma* involved in U insertion/deletion RNA editing. *RNA* **10**: 159-170.
- Sinha, K.M., Hines, J.C., Downey, N., and Ray, D.S. 2004. Mitochondrial DNA ligase in *Crithidia fasciculata*. *Proc. Natl. Acad. Sci. USA* **101**: 4361-4366.
- Smíd, O., Horáková, E., Vilímová, V., Hrdý, I., Cammack, R., Horváth, A., Lukeš, J., and Tachezy, J. 2006. Knockdowns of mitochondrial iron-sulfur cluster assembly proteins IscS and IscU down-regulate the active mitochondrion of procyclic *Trypanosoma brucei*. *J. Biol. Chem.* **281**: 28679-28686.
- Stuart, K. D., Schnauffer, A., Ernst, N.L., and Panigrahi, A. K. 2005. Complex management: RNA editing in trypanosomes. *Trends Biochem. Sci.* **30**: 97-105.
- Sturm, N. and Simpson, L. 1990. Kinetoplast DNA minicircles encode guide RNAs for editing of cytochrome oxidase subunit III mRNA. *Cell* **61**: 879-884

Vanhamme, L., Pérez-Morga, D., Marechal, C., Speijer, D., Lambert, L., Geuskens, M., Alexandre, S., Ismaili, N., Göringer, U., Benne, R., and Pays, E. 1998.

Trypanosoma brucei TBRGG1, a mitochondrial oligo(U)-binding protein that co-localizes with an *in vitro* RNA editing activity. *J. Biol. Chem.* **273**: 21825-21833.

Vassella, E., Kramer, R., Turner, C.M.R., Wankell, M., Modes, C., Van den Bogaard, M., and Boshart, M. 2001. Deletion of a novel protein kinase with PX and FYVE-related domains increases the rate of differentiation of *Trypanosoma brucei*. *Mol. Microbiol.* **41**: 33-46.

Vondrušková, E., van den Burg, J., Zíková, A., Ernst, N.L., Stuart, K., Benne, R., and Lukeš, J. 2005. RNA interference analyses suggest a transcript-specific regulatory role for MRP1 and MRP2 in RNA editing and other RNA processing in *Trypanosoma brucei*. *J. Biol. Chem.* **280**: 2429-2438.

Weng, J., Aphasizheva I., Etheridge, R. D., Huang L., Wang, X., Falick, A. M., Aphasizhev R. 2008. Guide RNA-binding complex from mitochondria of trypanosomatids. *Mol. Cell* **32**: 1-12.

Zíková, A., Kopečná, J., Schumacher, M.A., Stuart, K.D., Trantírek, L., and Lukeš, J. 2008. Structure and function of the native and recombinant mitochondrial MRP1/MRP2 complex from *Trypanosoma brucei*. *Int. J. Parasitol.* **38**: 901-912.

Figure legends

Fig. 1. Subunits of the MRB1 complex are essential for growth of procyclic stage trypanosomes. Cell densities (cells/ml) that were measured every 24 hours are plotted on a logarithmic scale on the y-axis over 14 days. Cells were diluted to 2×10^6 cells/ml every 2 days. Cells grown in the presence or absence of the RNAi-induction agent 1 μ g/ml tetracycline are indicated by grey (tet+) or black (tet-) lines, respectively. Growth curves of knockdowns of GAP1(**A**), GAP2 (**B**), RNA helicase (**C**) and Nudix hydrolase (**D**) are shown.

Fig. 2. The stability of the GAP proteins is dependent on their mutual association. (**A**) Crude mt lysates from 10^8 cells were loaded per lane on a SDS-PAGE gel and transferred onto a PVDF membrane for immunodecoration with either α -GAP1 or α -GAP2 polyclonal antibody, as indicated on the left. Antibody against the mitochondrial protein frataxin was used as a loading control. Lysates from cells grown for 3 days in the presence or absence of 1 μ g/ml tetracycline are indicated at

the top ("+" or "-") for either the GAP1 or GAP2 RNAi knockdown. 50 and 15 kDa protein markers are indicated on the right.

(B) Samples from the RNA helicase and Nudix hydrolase RNAi knockdown cells grown 4 days in the presence and absence of 1 μ g/ml tetracycline were prepared and labeled as described above.

Fig. 3. Effects of RNAi-silencing of the MRB1 complex subunits on maxicircle transcripts. Quantitative real time PCR analysis of pre-edited, edited and never-edited mRNAs and rRNAs was performed on knockdowns of GAP1 (A), GAP2 (B), RNA helicase (C) and Nudix hydrolase (D). The reaction for each amplicon was done in triplicate on cDNA generated from cells grown in the presence or absence of the RNAi-inducer tetracycline. The relative abundance of each of the examined RNAs upon the synthesis of interfering dsRNA was plotted on a logarithmic scale: 1 represents the wild-type levels, while a value above or below this number signifies an increase or decrease of a given RNA, respectively. These values were normalized to the cytosolic transcripts β -tubulin or 18S rRNA, whose levels are not affected by RNAi. The average (AVE $\sigma(C_T)$) and median (MED $\sigma(C_T)$) of the measured cycle threshold (C_T) values for all the reactions performed for each knockdown are indicated under each bar graph. The following pre-edited (P) and edited (E) RNAs were assayed: ATPase subunit 6 (A6), cytochrome oxidase subunits 2 (cox2) and 3 (cox3), cytochrome reductase subunit b (cyB), maxicircle unknown reading frame 2 (MURF2), NADH dehydrogenase subunit 7 (ND7) and ribosomal protein S12 (RPS12). The following never-edited RNAs were assayed: 9S RNA, 12S RNA, cox1, and ND4. The appropriate cytoplasmic mRNA targeted by RNAi were assayed for each knockdown, and are indicated on the right.

Fig. 4. Effects of RNAi-silencing of the MRB1 complex subunits on minicircle transcripts. The total population of minicircle encoded gRNAs were labeled in knockdowns of GAP1 (A), GAP2 (B), RNA helicase (C) and Nudix hydrolase (D). Labeling reactions were performed with guanylyltransferase, [α - 32 P] GTP and either 2.5 (A and B) or 5 μ g RNA (C and D). The gRNAs, appearing as a ladder mainly because of their heterogeneous 3'-oligo(U) tails, are indicated on the left. The top band marked with the "*" is a cytosolic RNA that is also labeled in this reaction and is used as a loading control.

Fig. 5. GAP1 assembly into macromolecular complexes is affected in the knockdowns for subunits of the MRB1 complex. Cleared lysates from hypotonically isolated mitochondria are separated by sedimentation in glycerol gradients. Odd fractions from the gradients were run on SDS-PAGE gels and transferred onto PVDF membranes.

(A) All gradients were verified by immunodecoration with monoclonal antibodies against editosome subunits KREPA2, KREL1 and KREPA3 (top panel), as well as Tbmp45 (bottom panel). A representative blot for GAP1 samples is shown here in which the appropriate antibody signals are indicated on the left. The presence (+) or absence (-) of tetracycline is specified on the right.

(B) α -GAP1 antibody was used to probe the glycerol gradient fractions from knockdown cell lines grown in the absence (-) or presence (+) of tetracycline as indicated on the right. The GAP1, GAP2, RNA helicase and Nudix hydrolase cell lines are indicated on the left.

Fig. 6. Evidence that mt RNA polymerase transcribes kDNA minicircles.

(A) Effect of RNAi induction on the mtRNAP mRNA was followed by Northern blot analysis. Its levels were analyzed by blotting 10 μ g of total RNA extracted from non-induced cells (-) and cells 3 days after RNAi induction (+) with a probe hybridizing to the gene fragment used for generation of dsRNA. The position of the targeted mRNA and the inducibly synthesized dsRNA are indicated by black and grey arrowheads, respectively. To visualize characteristic rRNA bands, used here as a loading control, the gel was stained with ethidium bromide.

(B) The same RNA was hybridized with a labeled probe for never-edited mRNA encoding cytochrome oxidase subunit 1 (cox1).

(C) Guanylyltransferase labeling of gRNAs using same RNA samples as used in (A) and (B) are performed and marked as in Fig. 4. 5 μ g of total RNA was used per lane.

Fig. 7. GAP1 assembly into large macromolecular complexes is reduced in cells with RNAi-silenced mt RNA polymerase. Western blot analysis of glycerol gradients was performed and marked as described in Fig. 5. Western blots of fractions were verified using the α -KREL1 (top panel) or α -Tbmp45 (bottom panel) monoclonal antibodies (A) or probed using the α -GAP1 polyclonal antibody (B).

Fig. 8. GAP RNAi are essential for gRNA biogenesis in bloodstream stage.

(A) Whole cell lysates from 1.5×10^7 cells were immunoblotted with either the α -GAP1 or α -GAP2 antibody for each of the GAP1 (top) and GAP2 (middle) cell lines. Cells grown in the absence (-) or presence (+) of 1 μ g/ml of tetracycline are indicated, as well as the days post-induction in the latter samples. An antibody decorating the cytosolic enolase was used as a loading control (bottom). All antibodies are indicated on the left.

(B) RNAi-silencing of the GAP proteins in the bloodstream stage results in growth inhibition. Growth curves over 7 days are shown for the GAP1 (top) and GAP2

(bottom) knockdown cell lines and are labeled as in Fig. 1. Cells were diluted every 24 hours to a density of 10^5 cells/ml.

(C) Guide RNA levels are diminished in the GAP-silenced bloodstream stage. The guanylyltransferase assay was performed on 5 μ g of total RNA as described in Fig. 4. The gels shown for GAP1 (left) and GAP2 (right) are also labeled as in Fig. 4., with the days post-induction by tetracycline indicated below the "+".

Fig. 9. The GAP proteins are localized in punctuate loci along the reticulated mitochondrion of procyclic *T. brucei*.

(A) Immunolocalization using rabbit α -GAP1 antibody in the GAP1 non-induced (top row, labeled Tet-) and RNAi-induced cells (bottom row, Tet+). Signal from cy3-conjugated α -rabbit antibody (Cy3), MitoTracker Green FM (Mito), the two images merged with DAPI staining of nucleus and kDNA (Merge-DAPI) and a composite image with a differential interference contrast image (Merge-DIC) are indicated along the top. White arrowheads indicate position of kDNA.


(B) Immunolocalization of the HA₃-tagged versions GAP1 (top row) and GAP2 (bottom row), which is immunodecorated with rabbit polyclonal antibody against the epitope. Signal from cy3-conjugated α -rabbit antibody (Cy3), DAPI staining of nucleus and kDNA (DAPI), a merged image of the two (Merge) and a composite image with a differential interference contrast image (Merge-DIC) are indicated along the top. White arrowheads indicate position of kDNA.

(C) Immunoblot analysis of the localization of overexpressed HA₃-tagged GAP1 (left) and 2 (right) proteins. Whole cell lysates (W) and digitonin extracted cytosolic (C) and mitochondrial (M) fractions of these cell lines grown for 2 days in the presence of 1 μ g/ml tetracycline. Antibody against the HA₃-tag was used as well as those against the MRP2 and enolase proteins, serving as mitochondrial and cytosolic markers, respectively. 10 μ g of protein of each digitonin fraction was loaded per lane.


Table 1. Nomenclature of examined MRB1 complex proteins.

Nomenclature introduced in this study	Nomenclature introduced by Weng et al. (2008)	GeneDB accession number
GAP1	GRBC1	Tb927.2.3800
GAP2	GRBC2	Tb927.7.2570
Nudix hydrolase	MERS1	Tb927.11.7290
RNA helicase	--	Tb927.4.1500

Fig. 1



A



B




Fig. 3









Fig. 4



A



B






Fig. 6



A



B








Fig. 8





Chapter 4:

Adaptations of *Trypanosoma brucei* to gradual loss of kinetoplast DNA: *Trypanosoma equiperdum* and *Trypanosoma evansi* are petite mutants of *T. brucei*.

(reprint of Lai *et al.*, 2008, Proc. Natl. Acad. Sci. USA 105, 1999-2004)

Abstract:

Trypanosoma brucei is a kinetoplastid flagellate, the agent of human sleeping sickness and ruminant nagana in Africa. Kinetoplastid flagellates contain their eponym kinetoplastDNA(kDNA), consisting of two types of interlocked circular DNA molecules: scores of maxicircles and thousands of minicircles. Maxicircles have typical mitochondrial genes, most of which are translatable only after RNA editing. Minicircles encode guide RNAs, required for decrypting the maxicircle transcripts. The life cycle of *T. brucei* involves a bloodstream stage (BS) in vertebrates and a procyclic stage (PS) in the tsetse fly vector. Partial [dyskinetoplastidy (Dk)] or total [akinetoplastidy (Ak)] loss of kDNA locks the trypanosome in the BS form. Transmission between vertebrates becomes mechanical without PS and tsetse mediation, allowing the parasite to spread outside the African tsetse belt. *Trypanosoma equiperdum* and *Trypanosoma evansi* are agents of dourine and surra, diseases of horses, camels, and water buffaloes. We have characterized representative strains of *T. equiperdum* and *T. evansi* by numerous molecular and classical parasitological approaches. We show that both species are actually strains of *T. brucei*, which lost part (Dk) or all (Ak) of their kDNA. These trypanosomes are not monophyletic clades and do not qualify for species status. They should be considered two subspecies, respectively *T. brucei equiperdum* and *T. brucei evansi*, which spontaneously arose recently. Dk/Ak trypanosomes may potentially emerge repeatedly from *T. brucei*.

Chapter 5:

The protein product of pan-edited ATP synthase subunit 6 is incorporated into a non-canonical Complex V in *Trypanosoma* species.

(submitted to Eukaryotic Cell)

The protein product of pan-edited ATP synthase subunit 6 is incorporated into a non-canonical complex V in *Trypanosoma* species

Hassan HASHIMI¹, Vladislava BENKOVIČOVÁ², Petra ČERMÁKOVÁ², De-Hua LAI¹, Anton HORVÁTH² and Julius LUKES^{1,*}

¹ Biology Centre, Institute of Parasitology, Czech Academy of Sciences, and Faculty of Biology, University of South Bohemia, České Budějovice (Budweis), Czech Republic; ² Department of Biochemistry, Faculty of Natural Sciences, Comenius University, Bratislava, Slovakia

Throughout eukaryotes, the gene encoding subunit 6 (ATP6) of the F₁F₀ ATP synthase (complex V) is maintained in mitochondrial genomes, presumably because of its high hydrophobicity due to its incorporation into the membrane-bound F₀ moiety. In *Trypanosoma* species, a mitochondrial (mt) transcript that undergoes extensive processing by RNA editing has a very low sequence similarity to ATP6 from other organisms. The notion that the putative ATP6 subunit is assembled into the F₀ sub-complex is ostensibly challenged by the existence of naturally occurring dyskinetoplastic (Dk) and akinetoplastid (Ak) trypanosomes, which are viable despite lacking the mt DNA required for its expression. Taking advantage of the differential phenotype between RNAi knockdown cell lines in which the expression of proteins involved in mitochondrial RNA metabolism and editing can be silenced, we provide several lines of support that ATP6 is encoded in the mt genome of *Trypanosoma* species and that it is incorporated into complex V. The reduction of the F₁F₀ oligomer of complex V coincides with the accumulation of the F₁ moiety in ATP6-lacking cells, which also appear to lack the F₀ ATP9 multimeric ring. The oligomycin sensitivity of ATPase activity of complex V in ATP6-lacking cells is reduced, reflecting the insensitivity of the Dk and Ak cells to this drug. In addition, the F₁ moiety of complex V appears to exist as a dimer in steady state conditions and contains the ATP4 subunit traditionally assigned to the F₀ sub-complex.

* To whom correspondence should be addressed: Institute of Parasitology, Branišovská 31, 37005 České Budějovice, Czech Republic. Tel.: 420-38-7775416; Fax: 420-38-5310388; E-mail: jula@paru.cas.cz

Keywords: RNA editing, ATP synthase, mitochondrion, *Trypanosoma*, respiratory complex, membrane potential.

INTRODUCTION

Trypanosoma brucei is a pathogen with a broad impact on the inhabitants of sub-Saharan Africa, subspecies of which are the causative agents of human sleeping sickness and ruminant nagana. The haemoflagellate switches between the tsetse fly vector and mammalian host, where it exists in the procyclic (PS) and bloodstream (BS) stages, respectively, and has to adapt to the different environments of the insect midgut and mammalian bloodstream. During its life cycle the parasite

undergoes substantial physiological and morphological changes [1], the transformation of its single mitochondrion belonging to the most dramatic of them [2]. The mitochondrion in the BS is considerably reduced and appears to lack cristae [reviewed in 3,4]. This morphological difference, in addition to the absence of cytochrome-containing respiratory complexes III and IV, reflects the exclusivity of glycolysis in the energy metabolism of *T. brucei* in the glucose-rich bloodstream of the mammalian host. In contrast, the PS contains a fully developed mitochondrion, appearing as a reticulated structure meandering throughout the cytoplasm, with the participation of complexes III and IV in the generation of ATP via oxidative phosphorylation [5].

Although the BS has a reduced mitochondrion, it is hardly a dormant organelle, as it still requires the expression of the mitochondrial (kinetoplast) DNA (kDNA), consisting of thousands of interlocked minicircles and dozens of maxicircles [reviewed in 6,7]. The latter molecules are the equivalents of classical mitochondrial (mt) DNA, containing genes required for the organelle's biogenesis, most of which encode subunits of the respiratory complexes. The expression of majority of these genes is not as straightforward as in other eukaryotes however, as their maturation requires extensive RNA editing in the form of post-transcriptional insertion and/or deletion of uridines (U) in preordained positions of the mRNA. A diverse population of minicircles encode small guide (g) RNAs, which direct the enzymatic machinery encompassed by the editosome protein complex to properly edit the mRNAs [reviewed in 8,9].

Trypanosomes maintain the mt membrane potential ($\Delta\Psi_m$) requisite for the import of essential proteins [10,11] and Ca^{2+} [12]. Typically, eukaryotes rely on respiratory complexes I, III and IV to pump protons outside of the mt matrix to generate $\Delta\Psi_m$, and the latter two complexes indeed have such a role in the PS [13]. However, noteworthy experiments have shown that in the BS it is the F_0F_1 -ATP synthase (complex V) that adopts the role of sustaining $\Delta\Psi_m$ [12,14,15].

In a typical mitochondrion, complex V acts as a dynamo driven by the flux of protons down the electrochemical gradient inherent in $\Delta\Psi_m$, which is coupled to the production of ATP. Its mechanism of action can be explained by exploring the remarkably conserved structure of complex V, the basic architecture of which is present in bacteria, archaea and eukaryotes [reviewed in 16,17]. Complex V is composed of two parts: the hydrophobic F_0 moiety embeds the complex into the mt inner membrane, while the F_1 sub-complex extends into the mt matrix. The F_0 part is composed of one ATP6 subunit (subunit *a* in the mammalian nomenclature), one or two ATP4 (*b*) subunits and a ring of 10-15 ATP9 (*c*) subunits. It is believed that this moiety mediates proton flux through the inner membrane via the interface between ATP6 and the ATP9-multimer ring, powering its rotary motion [17]. The resulting energy is used for ATP synthesis at the F_1 moiety, which is composed of a ring of three alternating α and β subunit pairs, a central stalk made up of the γ and ϵ subunits. The central stalk connects the spinning ATP9-composed ring to the stationary $\alpha_3\beta_3$ hexamer, facilitating conformational changes to the three catalytic sites present in the latter, which in turn allows ATP synthesis. Interaction between the F_0 ATP4 and F_1 OSCP subunits forms the peripheral stalk, which is believed to comprise part of a stator running parallel to the central stalk [18]. The action of complex V is reversed in the BS, in which ATP is consumed in order to pump protons out of the matrix [14,15].

The essential ATP6 subunit is retained in virtually all mt genomes, presumably due to its extreme hydrophobicity [19]. In *T. brucei*, a pan-edited mt transcript, requiring information from an estimated 21 gRNAs for the insertion of 448 Us and deletion of 28 Us to render a translatable ORF, has been hypothesized to encode ATP6 [20,21]. Such an assignment was supported by

analysis of the *in silico* translated product, revealing a similar hydropathy profile and low sequence similarity of the C-terminus to ATP6 orthologs from other species [20]. However, conclusive evidence for this speculative designation is missing [22].

The notion that this pan-edited mRNA encodes the ATP6 subunit, and that it is assembled into the F₀ moiety is, however, further complicated by the existence of natural and laboratory-induced dyskinetoplastic (Dk) and akinetoplastic (Ak) trypanosomes, which are viable despite a partial and complete loss of kDNA, respectively [reviewed in 23]. These cells are locked in the BS because they lack the mitochondrial-encoded subunits of the cytochrome-containing complexes III and IV, and are viable despite the absence of the ATP6 gene [23]. Yet RNA editing is essential in the BS of *T. brucei* [24], suggesting that the product of ATP6 is not entirely dispensable. Furthermore, the consumption of ATP by the F₁ moiety of complex V is required for maintaining $\Delta\Psi_m$, which is considered indispensable for the survival of any trypanosome, including the Dk and Ak strains [14]. One interpretation of these findings is that the protein product of the edited mRNA putatively assigned as ATP6 in the mt genome of *Trypanosoma* species is not incorporated into the F₀ moiety, perhaps because it does not encode this subunit. Another explanation postulates that in the Ak cells mutations have evolved in the nuclear-encoded subunits of the complex to offset the absence of ATP6, in which the ATPase activity of the F₁ is decoupled from the incomplete F₀ that lacks proton pumping capacity. A single amino acid substitution in the C-terminus of the γ subunit from a laboratory-induced Dk strain was elegantly demonstrated to be such a mutation [14], while additional mutations in the same region with a putative compensatory role were identified in several natural Ak strains [25]. Consistent with such a scenario is the insensitivity of *T. b. evansi* to oligomycin, a specific inhibitor of F₀, to which the BS *T. brucei* remains susceptible [14,26]. Furthermore, the putative ATP6 mRNA is edited in both PS and BS trypanosomes, while other transcripts are preferentially edited only in one of these stages [22,23].

Despite significant efforts in the last two decades, proteins translated from edited mRNAs remain elusive. So far, apocytochrome B (cyB) and cytochrome *c* oxidase subunit 2 (cox2) from the model trypanosomatid *Leishmania tarentolae* remain the only detected protein products of a moderately edited mRNA [27,28], whereas a protein translated from a pan-edited mRNA has yet to be found [28,29]. Recently, an important claim has been made that pan-editing generates alternative transcripts, leading to the production of several different proteins from one gene, a function analogous to alternative splicing [30]. Interestingly, only very few molecules were found to be fully edited and thus translatable into predicted subunits of respiratory complexes [31]. Such an inefficient production of these subunits, mostly deemed essential for a eukaryotic cell, further questions the presupposition that pan-editing creates translatable transcripts.

Unfortunately, direct recombinant manipulation of kDNA is still not feasible despite valiant efforts [32,33]. Thus, we have decided to address the translation and incorporation of the predicted ATP6 protein by taking advantage of the differential phenotypes resulting from RNA interference (RNAi)-mediated silencing of two nuclear-encoded proteins involved in mt RNA metabolism: mitochondrial RNA binding protein 2 (MRP2) and kinetoplastid RNA editing proteins KREPA6 (summarized in Table 1). MRP2 with its partner MRP1 forms a heterotetrameric MRP1/2 complex, which facilitates annealing of gRNAs to their cognate mRNAs [34]. RNAi knockdown (KDs) of MRP2 in the PS resulted in the downregulation of a subset of never-edited and edited mRNAs, with the important exception of ATP6 [35], which consequently caused the disruption of complexes III and IV (Table 1) [36]. Since KREPA6 is a subunit of the editosome [9], its depletion in the PS results in a general reduction of RNA

editing, affecting mRNAs for subunits of complexes III and IV, as well as ATP6 (Table 1) [37]. Although the secondary effect of its silencing on the respiratory complexes was not directly investigated, their disruption is considered inevitable, since this outcome is observed in the RNAi KD of the terminal uridylyl transferase required for gRNA maturation, which also affected all edited mRNAs in a similar fashion [38,39]. Here we present several lines of evidence, albeit indirect, for the existence of a genuine ATP6 subunit of an apparently non-canonical complex V in the mitochondrion of procyclic *T. brucei*.

EXPERIMENTAL PROCEDURES

Strains, cultivation and isolation of mitochondria - The MRP2 and KREPA6 RNAi cell lines were described elsewhere [35,37]. The PCR-generated fragment (forward primer with *Bam*HI restriction site: 5' - TGGGATCCAACACTGCACCATGGATTG - 3'; reverse primer with *Xho*I site: 5' - AAGCTCGAGTGGATGTTCTTTCCCTC - 3'; both restriction sites underlined) of ATP4 (p18) was cloned into the p2T7-177 vector. The ATP4 knockdown cell line was established by transformation of the parental 29-13 cell line with resulting cassette, as described elsewhere [40]. All transgenic cells, as well as the 29-13 cell line, were cultivated in SDM-79 medium under conditions described elsewhere [35]. The *T. brucei* strain 920, *T. b. equiperdum* strain 818 and *T. b. evansi* strain 810 were described previously [25]. The kinetoplast-mitochondrial vesicles from 5×10^8 non-induced and induced procyclic cells were isolated by hypotonic lysis as described elsewhere [13]. Pelleted mitochondrial vesicles were stored at -70 °C upon further use.

Quantitative real-time PCR - Quantitative real-time PCR reactions were performed as in Hashimi et al. [40]. Primers for amplification of pre-edited and edited mt mRNAs are as described elsewhere [41], as are those for KREPA6 [37]. The primer pair for detection of MRP2 cDNA is MRP2-qPCR-Fw (5' - GAAGCTTGGCTGTGTCCTTC-3') and MRP2-qPCR-Rv (5' - TGCGTCCGAATACGATTACA-3'). Relative RNA abundance between RNAi-induced and non-induced samples were calculated as described previously [25,37,40,41].

Measurement of ATPase activity and inhibition experiments- 1 mg of mt proteins was resuspended in 1 ml of the TC buffer (0,2 M KCl; 10 mM Tris-HCl, pH 8.2; 2 mM MgCl₂). The reaction was started at room temperature by adding ATP to the final concentration 5 mM and after 5 min was stopped by mixing 95 µl aliquots with 5 µl of 3 M CHCl₃-COOH, incubated for 30 min on ice, and spun (16 000 g for 10 min at 4°C). Ninety µl of the supernatant were added to 1 ml of the Sumner reagent (8,8% [w/v] FeSO₄; 375 mM H₂SO₄; 6,6% [w/v] (NH₄)Mo₇O₂₄). After 15 min incubation at room temperature, absorbance of free P_i was measured at 610 nm. Parallel experiments were performed with untreated samples and those in the presence of 10 µg/ml oligomycin or 1 mM azide. Inhibition experiments were reproduced in 10 independent measurements from four RNAi inductions with parallel non-induced controls. Statistical significance of observed differences in inhibition data were determined by Student's t-test.

In-gel activity staining and two-dimensional gel electrophoresis – For activity staining, 100 µg of mitochondrial lysate in 0.5 M aminocaproic acid and 2 % dodecylmaltoside was loaded per lane and analyzed on a 4-15% gradient blue native polyacrylamide (BN) PAGE gel as described

elsewhere [13]. Immediately after the run, the gel was transferred into either ATPase reaction buffer (35 mM Tris; 270 mM glycine; 19 mM MgSO₄; 0,3% [w/v] Pb(NO₃)₂; 11 mM ATP) for overnight incubation, or into cytochrome oxidase buffer (50 mM sodium phosphate, pH 7.4; 1 mg/ml 3,3'-diaminobenzidine; 24 U/ml catalase; 1 mg/ml cytochrome *c*; 75 mg/ml sucrose) for 3 hours staining, both by slow agitation. The ATPase activity appears as a white and cytochrome *c* oxidase activity as brown precipitate. The resulting bands were quantified with the scanning densitometry ImageQuant program (Molecular Dynamics). The BN PAGE gels for the first dimension of two-dimensional gel electrophoresis were prepared in the aforementioned fashion. The second dimension was resolved on 10% Tricine-SDS PAGE gels as in [13, 36].

Western blot analysis – After electrophoresis, the BN PAGE gel was blotted on a nitrocellulose membrane, and probed with polyclonal rabbit antibodies raised against complex V subunits (designation in parenthesis) of *Crithidia fasciculata* (F₁ moiety) [42], *Leishmania tarentolae* (ATP4) [43] and *T. brucei* (ATP9) [15] were used at 1:1,000, 1:3,000 and 1:500 dilutions, respectively. Secondary α -rabbit antibodies (1:2,000) (Sevapharma) coupled to horseradish peroxidase were visualized according to the manufacturer's protocol using the ECL kit (Pierce).

Isolation of F₁ moiety of complex V by chloroform extraction- In a protocol adapted from Linnet et al. [44], mitochondrial vesicles from 4 x 10⁸ cells were resuspended in 300 μ l STE buffer (0.25 M sucrose; 20 mM Tris.HCl, pH 7.9; 2 mM EDTA), and sonicated 4 x 10 s with 30 s intervals in between. The sonicate was centrifuged at 100,000 g for 30 min at 4 °C, and the pellet was then resuspended in 300 μ l STE. A 150 μ l volume of chloroform was added and the two phases were emulsified by vortexing for 10 s and spun in a microcentrifuge at maximum speed for 5 min at room temperature. The aqueous phase was transferred into a new centrifuge tube and ultracentrifuged at 100,000 g for 30 min at 25 °C. The proteins present in the supernatant were concentrated to 0.4 μ g / μ l using Microcon YM-30 centrifugation filters (Millipore), in which the solvent buffer was replaced with the resuspension solution (0.5 M aminocaproic acid; 0.25 % dodecylmaltoside). This sample was divided and stored overnight at either 0°C or room temperature to assay for the cold lability of the isolated F₁ moiety. About 5 μ g of extracted proteins were loaded per lane of the BN PAGE gel.

Electron microscopy – Cells were washed in 0.1 M phosphate-buffered saline (PBS) and fixed in 2.5% (v/v) glutaraldehyde in 0.1 M sodium cacodylate buffer, pH 7.4, for 1 h at 4 °C, then washed in 0.1 M PBS and post-fixed with 2% (w/v) OsO₄ in the same buffer for 1 h at room temperature. After dehydration in ethanol, the cells were embedded in PolyBed (Polysciences) and sectioned. Thin sections were stained with lead citrate and uranyl acetate, and examined in a JEOL JEM 1010 microscope. In order to estimate the number of cristae in mitochondria of the PS and BS of *T. brucei*, as well as KREPA6-silenced cells, the cristae were counted in over 200 carefully examined cross-sections of mitochondria from each cell line.

RESULTS

In-gel ATPase activity reveals oligomerization of complex V and dimerization of F₁ moiety in PS
- The lysates from hypotonically isolated mitochondria from the parental PS 29-13 cell line, from which the MRP2 and KREPA6 RNAi KD cells are derived, were resolved on a BN-PAGE gel.

Complex V was visualized by subsequent staining for in-gel ATPase activity (Fig. 1A), revealing two major upper ~2700 kDa and lower ~900 kDa bands. Both appear to be comprised of doublets or multiple smaller bands, with the bottom signal being the most intense. A smear is present between the upper and lower major bands, which are often resolved as multiple bands of less intensity than the upper and lower bands (see below). In order to confirm that these ATPase activity bands represent genuine complex V, Western blots of lysates separated by BN-PAGE were probed with two antibodies immunodecorating subunits of complex V. The α -subunit β and α -ATP4 antibodies, used in previous studies of *T. brucei* [14, 25], were originally raised against the F₁ moiety of *Crithidia fasciculata* [42], and the F₀-subunit of *Leishmania tarentolae* [39,43] (Figs. 1B and C). Both antibodies positively labeled all the bands with the same relative intensities as the in-gel activity stain. Polyclonal antibodies raised against the *Saccharomyces cerevisiae* ATP4, 6 and 9 subunits did not cross react with antigens present in *T. brucei* (data not shown).

Depending on the conditions used, multiple bands are resolved by native gel electrophoresis that correspond to different forms of complex V in yeast, such as oligomers, dimers and monomers of F₁F₀, as well as free F₁ particles [45]. We attempted to identify the form of complex V corresponding to the upper and lower bands, as well as the smear in between, by liquid chromatography-tandem mass spectroscopy (LC-MS/MS). However, only F₁ subunits were identified in all three samples, plus ATP4, possibly because the technique is not amenable to identification of the hydrophobic peptides comprising F₀ (data not shown). The presence of the canonical F₀ ATP4 in all bands also complicated this task. We resorted to the assignment of these bands by how their estimated molecular weights match the canonical stoichiometry [16, 17, 18] of the currently annotated complex V subunits in the GeneDB *T. brucei* genome database (<http://www.genedb.org/genedb/tryp/>). The top ~2700 kDa band has been provisionally assigned as an oligomer of complex V. The identity of the lower ~900 kDa doublet was more difficult to ascertain, since its migration could correspond to either a monomer of complex V or a dimer of the F₁ sub-complex. The isolated F₁ moiety from *L. tarentolae* has been reported to have a similar ~900 kDa size that we observe in *T. brucei*, and was also proposed to be a dimer [46].

To address this issue, the soluble F₁ moiety was prepared by chloroform extraction from hypotonically isolated mitochondria from the 29-13 cells following the protocol of Linnett et al. [44]. The F₁ moiety purified by this method from the membrane-bound hydrophobic subunits was electrophoresed on a BN gel along with the whole mitochondrial lysate. Upon staining by the in-gel ATPase activity assay, the lower band from the mt lysate migrates the same distance as the isolated F₁ moiety (Fig. 1A). The size implies the catalytic portion of complex V exists as a dimer, whose activity is cold labile (Fig. 1A), a typical characteristic of the isolated F₁ sector. Western analyses using antibodies against the F₁ β subunit and F₀ subunit ATP4 showed that both are present in the chloroform extracted F₁ sub-complex (Figs. 1B and C). ATP4 was also reported to be present in the chloroform extracted F₁ moiety of *C. fasciculata* [42].

ATP4 is required for the integrity of the F₁ moiety – The presence of ATP4, traditionally assigned to the F₀ moiety [16,17], in the chloroform-extracted F₁ sub-complex prompted us to generate RNAi KD of this subunit to investigate its position within the complex. Silencing of ATP4 resulted in slower growth of the procyclic cells (Fig. 2A). After 90 hours of RNAi-induction, the steady-state level ATP4 protein was virtually eliminated (Fig. 2A; inset), which was the time point chosen for subsequent experiments. In-gel ATPase activity and Western blot analysis using antibody against the β subunit showed that the catalytic portion of complex is

destabilized upon downregulation of ATP4 (Figs. 2B and C). This result is reminiscent of destabilization of the β subunit of F_1 upon silencing of its partner, the α subunit, and *vice versa* [14, 15], which may be an indication that ATP4 interacts directly with this moiety as well. This interpretation is consistent with its apparent association with isolated F_1 particle, as well as its interaction with the F_1 OSCP subunit of complex V. In any case, it is clear that ATP4 plays a role in maintaining the integrity of this moiety.

Differential effect of KREPA6 and MRP2 RNAi silencing on ATP6 editing - Inhibition of cell growth was re-evaluated in the PS cells, in which the KREPA6 or MRP2 transcripts were downregulated by RNAi (Figs. 3A and C). In agreement with previous reports [35,37], tetracycline induction of double stranded RNA targeting of these transcripts resulted in slower growth. Based on these results, subsequent experiments were performed on cells grown for six days in the presence and absence of tetracycline. This time point was chosen since our study concerns a secondary phenotype affecting the respiratory complexes, a strategy already employed in similar studies [36,39], taking advantage of the consequential results of the already direct effect of these RNAi KDs on mt RNAs.

In the original report, mt RNAs were assayed in the MRP2-silenced cells by poisoned primer extension [35], showing reduction of edited and never-edited subunits of complexes III and IV, without an apparent effect on the ATP6 mRNA (Table 1). We have verified the differential effect on ATP6 editing in the KREPA6 and MRP2 KDs using a more sensitive method based on quantitative real-time (q) PCR (Figs. 3B and D). Using previously described primers against pre-edited and edited ATP6 mRNAs [40], editing of this transcript is virtually abolished in the KREPA6 KD six days after tetracycline induction compared to its non-induced counterpart (Fig. 3B). In contrast, MRP2-silencing only slightly reduces the levels of pre-edited and edited ATP6 mRNAs (Fig. 3D), perhaps reflecting a general role in RNA stability and/or processing suggested for its eponymous complex [34,35]. Nonetheless, the level of the translatable ATP6 transcript is considerably closer to the wild-type level upon the depletion of MRP2 as compared to KREPA6.

A subset of the mt-encoded RNAs for subunits of respiratory complexes III and IV, plus the edited maxicircle unidentified reading frame 2 (MURF2), were also assayed by reverse transcription-qPCR to confirm that they are affected as originally reported in the KREPA6 and MRP2 KDs [35,37] (summarized in Table 1). As expected, the never-edited transcript of complex IV cytochrome *c* oxidase subunit 1 (*co1*) and the edited RNA of the complex III cytochrome reductase subunit b (*cyB*) were downregulated upon MRP2 silencing (Fig. 3D). Editing of *co3* and *cyB* was also reduced in the cells with downregulated KREPA6 (Fig. 3B). Moreover, editing of the MURF2 transcript, encoding a protein of unknown function, was equally affected in both KDs (Figs. 3B and D).

The accumulation of the F_1 moiety and reduction of the F_1F_0 oligomer of complex V in ATP6-depleted cells - Kinetoplasts from the non-induced and tetracycline-induced KREPA6 and MRP2 KDs were subjected to the aforementioned BN-PAGE/in-gel ATPase activity experiments. Complex V was visualized by running these lysates on the BN gels, followed by staining for the in-gel ATPase activity (Fig. 4A). In all the samples besides the KREPA6-silenced one, three bands between the upper and lower bands are present. The cells interfered against KREPA6, in which editing of ATP6 is virtually abolished (Fig. 3B), exhibit a reduction in oligomerization of complex V and a corresponding accumulation of free F_1 dimers (Fig. 4A). Semiquantification of

the ATPase bands reveal that the lowest doublet represents 96% of the sum density of the upper and lower bands in the lane containing the RNAi-silenced sample, while it comprises 57% and 58% in-gel activity from the untreated and parental cell lines (Fig. 4A). A similar enrichment of the free F_1 dimers was observed in KREN1 KD cells, in which one of the endonucleases essential for RNA editing has been ablated (data not shown) [47]. Such a disruption of complex V is not observed in the MRP2 KDs, in which editing of ATP6 persists, while the amount of never-edited, pre-edited and edited mRNAs of the other mitochondrial-encoded subunits of respiratory complexes is decreased [35,36]. The in-gel activity of free F_1 represents 32% and 67% of the sum density of the upper and lower bands observed in the non-induced and RNAi-induced samples, respectively (Fig. 4A). This pattern is apparent in western blot analyses with antibodies against the β and ATP4 subunits of the F_1 and F_0 sub-complexes, respectively (Figs. 4B and C). Interestingly, a signal corresponding to the uppermost of the three middle bands is observed in KREPA6-silenced lane of both immunodecorated blots. As expected, the in-gel activity of complex IV was dramatically reduced by about 70% when either MRP2 or KREPA6 are downregulated (Fig. 4D).

Formation of the F_0 ATP9 multimeric ring is disrupted in the absence of ATP6 – Two-dimensional gel electrophoresis was employed to further examine complex V in the KREPA6 KDs, since a polyclonal antibody against ATP9, a subunit forming the multimeric ring of the F_0 sub-complex, appears to recognize proteins resolved under denaturing conditions. Native complexes of RNAi cell lines grown in the presence and absence of tetracycline were separated on BN gels in the first dimension followed by SDS-PAGE in the second dimension to break apart individual subunits. As seen in Western blot in Fig. 5A, the β subunit and ATP4 have been detected in the upper and lower bands (as visualized by in-gel ATPase activity), as well as in the middle bands. However, ATP9 is confined to the region corresponding to the upper bands, and is also present to a lesser degree in the middle bands (Fig. 5A). In the KREPA6-silenced samples, ATP4 and F_1 are concentrated in the lowest bands, while ATP9 is undetectable, presumably because in the absence of ATP6, it cannot be assembled into the complex and is degraded (Fig. 5B). This finding also further supports the notion that the lower band represents a dimer of the F_1 moiety.

Sensitivity to oligomycin is decreased in ATP6-depleted kinetoplasts - The Ak trypanosome *T. b. evansi* is insensitive to oligomycin, a specific inhibitor of the F_0 subunit [14,26]. We investigated the possibility that the KREPA6-depleted cells may exhibit a decrease in sensitivity to oligomycin that would be reminiscent of the behavior of the ATP6-lacking Ak cells. The ATPase activity of hypotonically-isolated mitochondria from the non-induced and RNAi-induced KREPA6 and MRP2 KDs, as well as the parental 29-13 cells, were assayed for the release of P_i in the presence of 10 μ g/ml oligomycin and 1 mM azide, which inhibit the F_1 sub-complex (Fig. 6), as compared to samples without inhibitors. The observed ~40% reduction of ATPase activity by either inhibitor in mitochondria from the 29-13 cell is similar to what has been previously reported for the BS trypanosomes [14]. Downregulation of KREPA6 results in more than 50% lower sensitivity to oligomycin, as compared to the non-induced cells, while both cell lines do not differ in their sensitivity to azide (Fig. 6). The observed changes in oligomycin sensitivity are statistically significant ($p=0.0288$), while fluctuations in azide inhibition are not ($p=0.4907$), as determined by the Student's t-test. The MRP2 KD cells also exhibit somewhat reduced sensitivity to oligomycin, although they remain more susceptible to the drug than the KREPA6-

silenced cells. This relatively minor decrease in sensitivity of the MRP2-silenced cells was not statistically significant (oligomycin $p=0.1810$; azide $p=0.5199$), and may be due to the slight decrease of edited ATP6 mRNA, as revealed by qPCR (Fig. 3B).

Prevalence of the uncoupled F_1 moiety in Dk and Ak trypanosomes - In the induced KREPA6 KD cells, the accumulation of the F_1 moiety is apparent in the absence of the edited ATP6 mRNA (Fig. 4A). Therefore, in the BN-PAGE gels we have assayed the in-gel ATPase activity of lysed mitochondria purified from the Ak and Dk strains of *T. b. evansi* and *T. b. equiperdum*, respectively, as well as from the PS and BS of *T. brucei* (Fig. 7A). Two prominent bands are apparent in the BS that correspond to the top ~ 2700 kDa F_1F_0 -oligomer and the bottom ~ 900 kDa F_1 -dimer present in the PS cells. In the Dk and Ak strains only the lower ATPase activity band is apparent, however, with a higher intensity than that found in the BS cells (Fig. 7A). The BS and the Dk trypanosomes show considerably lower in-gel ATPase activity than the PS cells, which is in agreement with previous observations that complex V is downregulated in the stages infecting mammals [49]. The native forms of complex V were also visualized by Western blot analysis using an antibody directed against the β subunit, revealing the predominance of the F_1 dimer in the Dk and Ak trypanosomes (Fig. 7B). This finding was confirmed by Western blot analysis of two-dimensional gels resolving complex V, as in Fig. 5, showing that in the Ak *T. b. evansi* strain, the ATP4 subunit is restricted to the lower ATPase activity band.

Ultrastructure of PS mitochondrion is not altered in the absence of ATP6 - Oligomerization of complex V has been implicated in generating the inward folds of the mt inner membrane to form cristae [48]. Since multimers of complex V are reduced in the ATP6-depleted cells, we investigated this possibility by examining mitochondrial ultrastructure of the non-induced and induced KREPA6 and MRP2 KD cells by transmission electron microscopy. By this method, typical discoidal cristae of kinetoplastid flagellates were present in both KDs (Fig. 8A and data not shown), and their abundance was similar to that found in the PS parental 29-13 cells (Fig. 8B and Table 2). It is worth noting that cristae of very similar morphology were also observed in the BS of *T. brucei* (data not shown) and the Ak *T. b. evansi* (Fig. 8C), yet their abundance was about ten times lower in the mammalian stages (Table 2).

DISCUSSION

Since its seminal discovery [50], RNA editing in the mitochondrion of kinetoplastid flagellates is an intensely studied process [for review see 8,9]. There was very little doubt that the edited transcripts are translated into mt proteins. Still, so far only two proteins produced from a moderately edited and never-edited mt mRNA were shown to exist [27,28], while all attempts to detect predicted protein products of pan-edited mRNAs failed. An extensive analysis of mt transcripts in *T. brucei* and *L. tarentolae*, however, detected a widespread occurrence of partially edited molecules, with fully edited ones representing only a tiny fraction of the mt RNA population [51,52]. Yet evidence has recently been put forward that these are alternatively edited molecules translatable into different proteins [30,31]. In the particular case of pan-edited ATP6, only 10% of analyzed molecules could serve as blueprints for the synthesis of canonical ATP6 subunit of complex V [31].

Although perhaps inefficiently produced, the product of pan-edited ATP6 mRNA appears to be incorporated into the complex V of *Trypanosoma* species, as supported by several lines of evidence presented in this study. The PS cells, in which the 20S editosome subunit KREPA6 was RNAi-silenced, are deficient for ATP6, as compared to cells with downregulated MRP2. Indeed, RNA editing is generally inhibited by silencing proteins directly involved in the process, invariably affecting mt mRNAs encoding subunits of respiratory complexes, whereas complex V escaped such interference in the latter KD (Table 1 and Figs. 1B and D). The absence of the ATP6 protein caused by the disruption of RNA editing drastically reduces the level of F₁F₀ oligomers of complex V, while an accumulation of the free F₁ moiety, the soluble portion of the ATP synthase, takes place. The KREPA6 KD represents an ATP6-lacking background as compared to the MRP2-silenced cells, in which oligomerization occurs at approximately the wild type level. This situation is also apparent in Dk and Ak trypanosomes, which naturally lack the capacity to express ATP6, since they also exhibit an accumulation of the uncoupled F₁ moiety as compared to BS *T. brucei* containing the F₁F₀ oligomers. Furthermore, we show that ATP9, the subunit responsible for the formation of the multimeric ring of the F₀ moiety, is destabilized in the absence of ATP6. Finally, the sensitivity of complex V to oligomycin is reduced in cells lacking the capacity to edit ATP6 mRNAs, which correlates with the situation documented for the Dk trypanosomes naturally lacking ATP6 [14].

The deletion of ATP6 in *S. cerevisiae* resulted in the loss of F₁F₀ dimers and monomers and the appearance of free F₁, as revealed by in-gel ATPase activity staining [45]. Our findings in *T. brucei* are consistent with these results. However, ATP9 appears to be still assembled into complex V in the yeast null mutants [45]. In trypanosomes, the absence of ATP6 appears to destabilize the F₀ moiety, as implied by the diminished ATP9 ring, leading to the apparent deficiency of F₁F₀ formation. This observation indicates that ATP6 may not be the last subunit incorporated into complex V as has been suggested for the yeast ortholog, as a measure to prevent a premature leakage of protons by the incomplete sub-complex [45,53]. Furthermore, while complex V oligomerization has been demonstrated to participate in the formation of cristae in yeast [48], this does not appear to be the case in the ATP6-lacking *T. brucei*. Moreover, few cristae were even observed in the BS *T. brucei* and the Ak *T. b. evansi*, a surprising result given the paradigm that mitochondria of the mammalian stage do not contain these structures [2,3].

Several other differences between kinetoplastid and canonical complex V were underscored by this study. It appears that ATP4, a subunit traditionally assigned to F₀, is firmly associated with F₁ in these flagellates. This conclusion is viable given its role in the formation of the peripheral stalk by direct interaction with the F₁ OSCP subunit and is supported by several lines of evidence presented in this study. First, ATP4 is a constant presence within the soluble moiety, even after separation from the membrane-bound subunits by chloroform extraction [44], as also reported in *C. fasciculata* [42]. This localization is in contrast to ATP9, which is absent in the free F₁ moiety. Second, RNAi silencing of ATP4 leads to the destabilization of F₁, an indication of the mutual interdependence of interaction partners [13, 35] and reminiscent of studies showing that silencing of one of the subunits forming the F₁ αβ heterodimer results in the degradation of the other [14, 15]. In addition, silencing of the α subunit leads to degradation of ATP4 [15]. This phenotype is in contrast with the one resulting from the disruption of the ATP4 gene in *S. cerevisiae*, triggering the decoupling of F₀ from F₁ without destabilizing proteins of the latter moiety [54].

This study also suggests that the F₁ subunit exists as a dimer across kinetoplastid flagellates, as originally reported for *L. tarentolae* [46]. Dimerization of this moiety has been

reported in other eukaryotes [55]. However, in these organisms it is promoted by an inhibitor protein in response to conditions that lead to diminished $\Delta\Psi_m$, such as oxygen deprivation, in order to conserve ATP that would otherwise be spent by reversal of complex V [55]. The chloroform-extracted F_1 moiety of *T. brucei*, whose migration in native gels also suggests a dimer conformation, is apparently not inhibited, since they maintain in-gel ATPase activity. Thus, its apparent dimerization is formed by another mechanism. Homodimers of the ATP4 subunit, assigned to the membrane-bound F_0 sub-complex, have been demonstrated to form between two adjacent complexes in yeast [56]. The presence of this subunit in the isolated F_1 sector in trypanosomes may mediate the dimerization of this moiety. Nonetheless, the occurrence of F_1 as a dimer appears to be unique to these organisms.

Although the presented data suggest for the first time that a protein product of a pan-edited RNA is indeed translated, and in this case assembled into complex V, due to the limitations of our system, we were unable to directly address the question whether ATP6 is essential for ATP synthesis. A logical experiment would be to test *in vitro* ATP synthase activity [57] from mitochondria isolated from the analyzed KDs. However, this assay requires that the capacity to generate $\Delta\Psi_m$ is preserved, which both KDs lack due to the inevitable disruption of proton pumping complexes III and IV. Nevertheless, the notion that ATP6 is indispensable is further supported by reports that RNA editing is essential in the BS *T. brucei*, which uses complex V to sustain $\Delta\Psi_m$ [23], while its maturation by this process is redundant in the Dk and Ak trypanosomes because of compensatory mutations in some of the F_1 subunits [14,25].

Acknowledgements- We thank Salvador Tarun, Yuko Ogata and Ken Stuart (Seattle Biomedical Research Institute) for the gift of the KREPA6 and KREN1 cell lines and for facilitating mass spectroscopy analysis. We also thank Noreen Williams (State University of New York, Buffalo), Larry Simpson (University of California, Los Angeles), Jean Velours (Institut de Biochimie et Génétique Cellulaires, Bordeaux), Rob Benne and Dave Speijer (University of Amsterdam) for kindly providing antibodies. We thank Alena Ziková and Zdeněk Verner for fruitful discussions and Achim Schnauffer (University of Edinburgh) for critical reading of the manuscript. This work was supported by Grant Agency of the Czech Republic (204/06/1558), Grant Agency of the Czech Academy of Sciences (A500960705), and the Czech Ministry of Education (LC07032, 2B06129 and 6007665801) and Grant Agency of the Slovak Ministry of Education and the Slovak Academy of Sciences 1/3241/06.

REFERENCES

- 1 Matthews, K.R. (2005) The developmental cell biology of *Trypanosoma brucei* J. Cell Sci. **118**, 283-29
- 2 Vickerman, K. (1985) Developmental cycles and biology of pathogenic trypanosomes. Med. Bull. **41**, 105-114
- 3 Schneider, A. (2001) Unique aspects of mitochondrial biogenesis in trypanosomatids. Int. J. Parasitol. **31**, 1403–1415
- 4 Hannaert, V., Bringaud, F., Opperdoes, F.R., and Michels, P.A. M. (2003) Evolution of energy metabolism and its compartmentation in kinetoplastida. Kinetopl. Biol. Dis. **2**, 11–42
- 5 Besteiro, S., Barrett, M.P., Riviere, L., and Bringaud, F. (2005) Energy generation in insect stages of *Trypanosoma brucei*: metabolism in flux. Trends Parasitol. **21**, 185–191

- 6 Shlomai, J. (2004) The structure and replication of kinetoplast DNA. *Curr. Mol. Med.* **4**, 623–647
- 7 Liu, B., Liu, Y., Motyka, S.A., Agbo, E.E.C., and Englund, P.T. (2005) Fellowship of the rings: the replication of kinetoplast DNA. *Trends Parasitol.* **21**, 363–369
- 8 Lukeš, J., Hashimi, H., and Ziková, A. (2005) Unexplained complexity of the mitochondrial genome and transcriptome in kinetoplastid flagellates. *Curr. Genet.* **48**, 277-299
- 9 Stuart, K.D., Schnauffer, A., Ernst, N.L., and Panigrahi, A.K. (2005) Complex management: RNA editing in trypanosomes. *Trends Biochem. Sci.* **30**, 97–105
- 10 Nolan, D.P., and Voorheis, H.P. (1992) The mitochondrion in blood-stream forms of *Trypanosoma brucei* is energized by the electrogenic pumping of protons catalyzed by the F₁F₀-ATPase. *Eur. J. Biochem.* **209**, 207–216
- 11 Bertrand, K.I., and Hajduk, S.L. (2000) Import of a constitutively expressed protein into mitochondria from procyclic and bloodstream forms of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **106**, 249-260
- 12 Vercesi, A.E., Docampo, R., and Moreno, S.N. (1992) Energization-dependent Ca²⁺ accumulation in *Trypanosoma brucei* bloodstream and procyclic trypomastigotes mitochondria. *Mol. Biochem. Parasitol.* **56**, 251-7
- 13 Horváth, A., Horáková, E., Dunajčíková, P., Verner, Z., Pravdová, E., Šlapetová, I., Cuninková, L., and Lukeš J. (2005) Down-regulation of the nuclear-encoded subunits of the complexes III and IV disrupts their respective complexes but not complex I in procyclic *Trypanosoma brucei*. *Mol. Microbiol.* **58**, 116-130
- 14 Schnauffer, A., Clark-Walker, G.D., Steinberg, A.G., and Stuart, K. (2005) The F₁-ATP synthase complex in bloodstream stage trypanosomes has an unusual and essential function. *EMBO J.* **24**, 4029–4040
- 15 Brown, S.V., Hosking, P., Li, J., and Williams, N. (2006) ATP synthase is responsible for maintaining mitochondrial membrane potential in bloodstream form *Trypanosoma brucei*. *Eukaryot. Cell* **5**, 45–53
- 16 Boyer, P.D. (1997) The ATP synthase – a splendid molecular machine. *Ann. Rev. Biochem.* **66**, 717-749
- 17 Velours, J., and Arselin, G. (2000) The *Saccharomyces cerevisiae* ATP synthase. *J. Bioenerg. Biomemb.* **32**, 383-390
- 18 Mulkidjanian, A.Y., Makarova, K.S., Galperin, M.Y. and Koonin, E.V. (2007) Inventing the dynamo machine: The evolution of the F-type and V-type ATPases. *Nature Rev. Microbiol.* **5**, 892-899
- 19 Funes, S., Davidson, E., Claros, M.G., van Lis, R., Perez-Martinez, X., Vazquez-Acevedo, M., King, M.P., and Gonzales-Halphen D. (2002) The typically mitochondrial DNA-encoded ATP6 subunit of the F₁F₀-ATPase is encoded by a nuclear gene in *Chlamydomonas reinhardtii*. *J. Biol. Chem.* **277**, 6051-6058
- 20 Bhat, G.J., Koslowsky, D.J., Feagin, J.E., Smiley, B.L., and Stuart, K. (1990) An extensively edited mitochondrial transcript in kinetoplastids encodes a protein homologous to ATPase subunit -6. *Cell* **61**, 885–894
- 21 Corell, R.A., Feagin, J.E., Riley, G.R., Strickland, T., Guderian, J.A., Myler, P.J., and Stuart, K. (1993) *Trypanosoma brucei* minicircles encode multiple guide RNAs which can direct editing of extensively overlapping sequences. *Nucl. Acids Res.* **21**, 4313–4320

- 22 Stuart, K.D., Allen, T.E., Heidmann, S., and Seifert, S.D. (1997) RNA editing in kinetoplastid protozoa. *Microbiol. Mol. Biol. Rev.* **61**, 105–120
- 23 Schnauffer, A., Domingo, G.J., and Stuart, K. (2002). Natural and induced dyskinetoplastic trypanosomatids: how to live without mitochondrial DNA. *Int. J. Parasitol.* **32**, 1071–1084
- 24 Schnauffer, A., Panigrahi, A.K., Panicucci, B., Igo, R.P.Jr., Wirtz, E., Salavati, R., and Start, K. (2001) An RNA ligase essential for RNA editing and survival of the bloodstream form of *Trypanosoma brucei*. *Science* **291**, 2159-2162
- 25 Lai, D.-H., Hashimi, H., Lun, Z.-R., Ayala, F.J., and Lukeš, J. (2008) Adaptation of *Trypanosoma brucei* to gradual loss of kinetoplast DNA: *T. equiperdum* and *T. evansi* are petite mutants of *T. brucei*. *Proc. Natl. Acad. Sci. USA* **105**, 1999-2004
- 26 Opperdoes, F.R., Borst, P. and Derijcke, D. (1976) Oligomycin sensitivity of the mitochondrial ATPase as a marker for fly transmissibility and the presence of functional kinetoplast DNA in African trypanosomes. *Comp. Biochem. Physiol. B-Biochem. Mol. Biol.* **55**, 25-30
- 27 Horváth, A., Berry, E.A., and Maslov, D.A. (2000) Translation of the edited mRNA for cytochrome b in trypanosome mitochondria. *Science* **287**, 1639–1640
- 28 Horváth, A., Neboháčová, M., Lukeš, J. and Maslov, D.A. (2002) Unusual polypeptide synthesis in the kinetoplast-mitochondria from *Leishmania tarentolae* – Identification of individual de novo translation products. *J. Biol. Chem.* **277**, 7222-7230.
- 29 Panigrahi, A.K., Ogata, Y., Zíková, A., Anupama, A., Dalley, R.A., Acestor, N., Myler, P.J., and Stuart, K.D. (2009) A comprehensive analysis of *Trypanosoma brucei* mitochondrial proteome. *Proteomics* **9**, 434-50.
- 30 Ochsenreiter, T. and Hajduk, S.L. (2006) Alternative editing of cytochrome c oxidase III mRNA in trypanosome mitochondria generates protein diversity. *EMBO Rep.* **7**, 1128-1133
- 31 Ochsenreiter, T., Cipriano, M. and Hajduk, S.L. (2008) Alternative mRNA editing in trypanosomes is extensive and may contribute to mitochondrial protein diversity. *PLoS One* **3**, e1566
- 32 Sbicego, S., Schnauffer, A., and Blum B. (1998) Transient and stable transfection of *Leishmania* by particle bombardment. *Mol. Biochem. Parasitol.* **94**, 123-126
- 33 Schnauffer, A., Sbicego, S., and Blum B. (2000) Antimycin A resistance in a mutant *Leishmania tarentolae* strain is correlated to a point mutation in the mitochondrial apocytochrome *b* gene. *Curr. Genet.* **37**, 234-241
- 34 Schumacher, M.A., Karamooz, E., Zíková, A., Trantírek, L., and Lukeš, J. (2006) Crystal structures of *Trypanosoma brucei* MRP1/MRP2 guide-RNA-binding complex reveals RNA matchmaking mechanism. *Cell* **126**, 701-711
- 35 Vondrušková, E., van den Burg, J., Zíková, A., Ernst, N.L., Stuart, K., Benne, R., and Lukeš J. (2005) RNA interference analyses suggest a transcript-specific regulatory role for MRP1 and MRP2 in RNA editing and other RNA processing in *Trypanosoma brucei*. *J. Biol. Chem.* **280**, 2429-2438
- 36 Zíková, A., Horáková, E., Jirků, M., Dunajčíková, P., and Lukeš, J. (2006) The effect of down-regulation of mitochondrial RNA-binding proteins MRP1 and MRP2 on respiratory complexes in procyclic *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **149**, 65-73

- 37 Tarun, S.Z. Jr., Schnaufer, A., Ernst, N.L., Proff, R., Deng, J., Hol, W., and Stuart, K. (2008) KREPA6 is an RNA-binding protein essential for editosome integrity and survival of *Trypanosoma brucei*. *RNA* **14**, 347-358
- 38 Aphasizhev, R., Sbicego, S., Peris, M., Jang, S.-H., Aphasizheva, I., Simpson, A.M., Rivlin, A., and Simpson, L. (2002) Trypanosome mitochondrial 3' terminal uridylyl transferase (TUTase): The key enzyme in U-insertion/deletion RNA editing. *Cell* **108**, 637-648
- 39 Nebohacova, M., Maslov, D.A., Falick, A.M., and Simpson, L. (2004) The effect of RNA interference down-regulation of RNA editing 3'-terminal uridylyl transferase (TUTase) 1 on mitochondrial de novo protein synthesis and stability of respiratory complexes in *Trypanosoma brucei*. *J. Biol. Chem.* **279**, 7819-7825
- 40 Hashimi, H., Zíková, A., Panigrahi, A.K., Stuart, K.D., and Lukeš, J. (2008) TbRGG1, a component of a novel multi-protein complex involved in kinetoplastid RNA editing. *RNA* **14**, 970-980
- 41 Carnes, J., Trotter, J.R., Ernst, N.L., Steinberg, A., and Stuart, K. (2005) An essential RNase III insertion editing endonuclease in *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. USA* **102**, 16614-16619
- 42 Speijer, D., Breek, C.K., Muijsers, A.O., Hartly, A.F., Berden, J.A., Albrecht, S.P., Samyn, B., Van Beeumen, J., and Benne, R. (1997) Characterization of the respiratory chain from cultured *Crithidia fasciculata*. *Mol. Biochem. Parasitol.*, **85**, 171-186
- 43 Bringaud, F., Peris, M., Zen, K.H., and Simpson, L. (1995) Characterization of two nuclear-encoded protein components of mitochondrial ribonucleoprotein complexes from *Leishmania tarentolae*. *Mol. Biochem. Parasitol.* **71**, 65-79
- 44 Linnett, P.E., Mitchell, A.D., Partis, M.D., and Beechey, R.B. (1979) Preparation of the soluble ATPase from mitochondria, chloroplasts, and bacteria by the chloroform technique. *Methods Enzymol.* **55**, 337-343
- 45 Rak, M., Tetaud, E., Godard, F., Sagot, I., Salin, B., Duvezin-Caubet, S., Slonimski, P.P., Rytka, J., and di Rago, J.-P. (2007) Yeast cells lacking the mitochondrial gene encoding the ATP synthase subunit 6 exhibit a selective loss of complex IV and unusual mitochondrial morphology. *J. Biol. Chem.* **282**, 10853-10864
- 46 Nelson, R.E., Aphasizheva, I., Falick, A.M., Nebohacova, M., and Simpson, L. (2004) The I-complex in *Leishmania tarentolae* is a uniquely-structured F₁-ATPase. *Mol. Biochem. Parasitol.* **135**, 219-222
- 47 Trotter, J.R., Ernst, N.L., Carnes, J., Panicucci, B., and Stuart, K. (2005) A deletion site editing endonuclease in *Trypanosoma brucei*. *Mol. Cell* **20**, 403-412
- 48 Paumard, P., Vaillier, J., Coulary, B., Schaeffer, J., Soubannier, V., Mueller, D.M., Brethes, D., di Rago, J.P., and Velours, J. (2002) The ATP synthase is involved in generating mitochondrial cristae morphology. *EMBO J.* **21**, 221-230
- 49 Brown, S.V., Chi, T.B., and Williams, N. (2001) The *Trypanosoma brucei* mitochondrial ATP synthase is developmentally regulated at the level of transcript stability. *Mol. Biochem. Parasitol.* **115**, 177-187
- 50 Benne, R., Van den Burg, J., Brakenhoff, J.P., Sloof, P., Van Boom, J.H., and Tromp, M.C. (1986) Major transcript of the frameshifted *coxII* gene from trypanosome mitochondria contains four nucleotides that are not encoded in the DNA. *Cell* **46**, 819-826

- 51 Decker, C.J., and Sollner-Webb, B. (1990) RNA editing involves indiscriminate U changes throughout precisely defined editing domains. *Cell* **61**, 1001-1011
- 52 Sturm, N.R., and Simpson, L. (1990) Partially edited mRNAs for cytochrome b and subunit III of cytochrome oxidase from *Leishmania tarentolae* mitochondria: RNA editing intermediates. *Cell* **61**, 871-878
- 53 Tzagoloff, A., Barrientos, A., Neupert, W., and Hermann, J.M. (2004) Atp10p assists assembly of atp6p into the F₀ unit of the yeast mitochondrial ATPase. *J. Biol. Chem.* **279**, 19775-19780
- 54 Paul, M.F., Velours, J., Arselin de Chateaubodeau, G., Aigle, M., and Guerin B. (1989) The role of subunit 4, a nuclear-encoded protein of the F₀ sector of yeast mitochondrial ATP synthase, in the assembly of the whole complex. *Eur. J. Biochem.* **185**, 163-71.
- 55 Dienhart, M., Pfeiffer, K., Schagger, H., and Stuart, R.A. (2002) Formation of the yeast F₁F₀-ATP synthase dimeric complex does not require the ATPase inhibitor protein, Inh1. *J. Biol. Chem.* **277**, 39289-39295
- 56 Spannagel, C., Vaillier, J., Arselin, G., Graves, P.-V., Grandier-Vazeille, X., and Velours, J. (1998) Evidence of a subunit 4 (subunit b) dimer in favor of the proximity of ATP synthase complexes in yeast inner mitochondrial membrane. *Biochim. Biophys. Acta* **1414**, 260-264
- 57 Allemann, N., and Schneider, A. (2000) ATP production in isolated mitochondria of procyclic *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **111**, 87-94

FIGURE LEGENDS AND TABLES

Fig. 1. Oligomerization of complex V and dimerization of free F₁ moiety in procyclic *T. brucei*. 100 µg of protein from lysates of hypotonically isolated mitochondria and the equivalent of 5 µg of the F₁ moiety after chloroform extraction were electrophoresed on a 4-15% blue native gel. The position of the ferritin dimer (Sigma) used as molecular weight marker are indicated on the left. 0°C, F₁ moiety isolated by chloroform extraction and incubated overnight at 0°C; Chl, F₁ moiety isolated by chloroform extraction and incubated overnight at room temperature; Mt, lysates from hypotonically isolated mitochondria; u – upper activity bands (~2700 kDa); l – lower activity bands (~900 kDa).

After the run, the gel was either stained for in-gel ATPase activity (**A**), or transferred onto a Nitrocellulose membrane and immunodecorated with polyclonal antibodies against either the F₁ β subunit (**B**) or the F₀ subunit ATP4 (**C**).

Fig. 2. Downregulation of ATP4 results in slower growth and disassembly of the F₁ moiety. (**A**) The growth curves for non-induced (grey open circles) and RNAi-induced cells (black filled squares) are shown for a period of 14 days. The \bar{y} axis is depicted on a log scale and represents the cell densities of the cultures measured every 24 hours, as specified on the \bar{x} axis. Inset displays the downregulation of the ATP4 protein after 90 hours of RNAi-induction (+) as compared to the non-induced cells (-), as determined by Western blot analysis (Ab). Ponceau staining of the membrane is shown as a loading control (Pon).

(**B**) Complex V from mitochondrial lysates of the 29-13 parental cell line, as well as the ATP4 RNAi knockdown cell line grown for 90 hours in the absence (-) and presence (+) of tetracycline,

was resolved on a 4-15% blue native gel and visualized by the in-gel ATPase activity assay, as in Fig 1A. The upper (u, ~2700 kDa) and lower (l, ~900 kDa) bands, in addition to the position of the ferritin dimer marker (880 kDa), are indicated on the left and right, respectively. 100 μ g of protein was loaded per lane.

(C) Visualization of $F_1\beta$ subunit after the blue native gel was transferred to nitrocellulose and probed with specific antibody. Labeling as in Fig. 2B.

Fig. 3. Effect of RNAi-silencing of KREPA6 and MRP2 on cell growth and editing of ATP6 mRNA.

The growth curves for non-induced (grey open circles) and induced KREPA6 (A) and MRP2 (C) RNAi knockdowns (black filled squares) are shown for a period of 10 days and depicted as in Fig. 2A. Real time quantitative PCR analysis of a subset of mitochondrial-encoded RNAs, including pre-edited and edited ATP6 mRNAs, as well as the levels of the particular nuclear-encoded and RNAi-targeted mRNA (KREPA6 – B; MRP2 – D) was performed in triplicate on cDNAs generated from cells grown for 6 days in the presence or absence of tetracycline. For each target amplicon, the relative change in RNA abundance due to induction of RNAi-silencing of the particular transcript was determined by using cytosolic transcripts of β -tubulin (striped bar) and 18S rRNA (grey bar) as internal references, since their transcription was not affected by treatment. The following pre-edited (P) and edited (E) mRNAs were assayed: ATPase subunit 6 (A6), cytochrome oxidase subunits 1 (co1) and 3 (co3), cytochrome reductase subunit b (cyB) and maxicircle unknown reading frame 2 (MURF2). The average and median standard deviation of the measured triplicate cycle threshold (C_t) values are 0.13 and 0.10, respectively.

Fig. 4. Decrease of complex V F_1F_0 oligomers and accumulation of free F_1 dimers in the ATP6-deficient cells. Mitochondrial complexes from KREPA6 and MRP2 knockdown cells, grown for six days in the absence (-) or presence (+) of 1 μ g/ml tetracycline (tet), respectively, were separated by blue native gel electrophoresis and stained for ATPase activity as described in Fig. 1. 100 μ g of protein was loaded per lane. 29-13, procyclic *T. brucei* strain 29-13. The position of the ferritin dimer molecular weight marker is indicated on the left; u – upper activity bands (~2700 kDa); l – lower activity bands (~900 kDa).

(A) In-gel ATPase activity. The % density of the lower activity bands relative to that of total density of the combined bands is indicated at the bottom.

(B) Western blot analysis with antibody against the $F_1\beta$ subunit under the conditions described in Fig. 1.

(C) Western blot analysis with antibody against the F_0 subunit ATP4 under the conditions described in Fig. 1.

(D) In-gel cytochrome *c* oxidase (complex IV) activity. The presence of complex IV on blue native gels, resolved as described in Fig. 1, was detected by incubation with the reaction buffer (see Experimental Procedures), which yields a dark precipitate upon catalysis by the complex. In both RNAi-silenced cells, the density of the activity band is reduced by ~70% compared to their non-induced counterparts.

Fig 5. Incorporation of F₀ ATP9 into complex V in the presence and absence of ATP6. Complex V from KREPA6 knockdown cell lines grown for six days in the absence (A) and presence (B) of 1 µg/ml tetracycline, were resolved by two-dimensional gel electrophoresis. Native mitochondrial complexes were separated on a 4-15% blue native gel (1st D [first dimension]; top panel), as described in Fig. 1. Complex V was subsequently visualized by in-gel ATPase activity and the upper ~2700 kDa (u) and lower ~900 kDa (l) bands are indicated just below. The individual subunits of the complexes were separated by 10% Tricine SDS-PAGE (2nd D [second dimension], three bottom panels), transferred to nitrocellulose and probed with antisera against the F₁ β-subunit, ATP4 and ATP9. These antibodies are indicated on the left. Molecular weight markers are specified on the right.

Fig. 6. ATP6-depleted cells exhibit reduced sensitivity to the F₀ inhibitor oligomycin. Inhibition of the *in vitro* ATPase activity of complex V by 10 µg/ml of oligomycin (grey bar), which targets the F₀ moiety, and 1 mM azide (striped bar), interfering with the F₁ moiety, was assayed. Hypotonically isolated mitochondria from the parental 29-13 cell line and from the KREPA6 and MRP2 knock-down cells, grown for six days in the absence (-) and presence (+) of 1 µg/ml tetracycline, were solubilized and incubated with 5 mM ATP. The release of free P_i was measured by absorbance at 610 nm. The vertical axis depicts average % inhibition of ATPase as compared to the untreated samples. For both inhibitors, three independent experiments using mitochondria from the 29-13 cells and six each from individual RNAi-induced and non-induced KREPA6 and MRP2 KDs were performed. Error bars indicate standard deviation among the ten individual experiments. The statistical significance of % inhibition of ATPase activity by oligomycin was p= 0.0288 and p= 0.1810 in between induced (+) and non-induced (-) KREPA6 and MRP2 KD cells, respectively, as determined by Student's t-test. P-values for changes in sensitivity to azide inhibition were p= 0.4907 and p= 0.5199 for KREPA6 and MRP2 tet+ and – samples.

Fig. 7. The prevalence of the F₁ moiety in naturally akinetoplastic *T. b. evansi* and dyskinetoplastic *T. b. equiperdum*.

(A) In-gel ATPase activity. Mitochondrial protein complexes were resolved and stained as described in Fig. 1. 100 µg of protein was loaded per lane. BS, bloodstream *T. brucei*, strain 920; PS, procyclic *T. brucei* strain 29-13; Ak, akinetoplastic bloodstream *T. brucei evansi*, strain 810; Dk, dyskinetoplastic bloodstream *T. brucei equiperdum*, strain 818; u – upper activity band (~2700 kDa); l – lower activity bands (~900 kDa).

(B) Western blot analysis with antibody against the F₁ β subunit under the conditions described in Fig. 2. The position of the ferritin dimer molecular weight marker is indicated on the left.

(C) Two-dimensional gel electrophoresis resolution of complex V in the bloodstream *T. brucei* (BS) was performed and labeled as described in detail in Fig. 5.

(D) Two-dimensional gel electrophoresis resolution of complex V in the akinetoplastic bloodstream *T. brucei evansi*, strain 810 (Ak) was performed and labeled as described in detail in Fig. 5.

Fig. 8. The mitochondrion in the ATP6-deficient cells has unaltered morphology and does not lose the cristae. Transmission electron micrographs taken from the KREPA6 knock-down cells grown for six days in the presence (+) of 1 µg/ml tetracycline (**A**), parental 29-13 cell line (procyclic *T. brucei*) (**B**), and the bloodstream stages of naturally akinetoplastic *T. b. evansi* (**C**). Typical discoidal cristae are labeled with arrows; n = nucleus; k = kinetoplast; f = flagellum. Bar – 500 nm (**A**), 200 nm (**B**) and 1 µm (**C**).

Table 1. Summary of mitochondrial transcripts affected by the RNAi silencing of the KREPA6 and MRP2 proteins.

RNAi KD	Effect on mitochondrial RNAs						Reference
	I	III	IV	rRNA	RPS12	ATP6	
KREPA6	E	E	E	ND	E	E	35
MRP2	NE & E	E	NE	0	E	0	33

Mitochondrial RNAs: I = subunit of complex I (NADH dehydrogenase); III = subunit of complex III (cytochrome *c* reductase); IV = subunit of complex IV (cytochrome *c* oxidase); rRNA = mitoribosomal RNA; RPS12 = ribosomal protein subunit 12.

Effect: E = decrease in edited mRNA; NE = decrease in never-edited RNA; 0 = no effect; ND = not determined.

Table 2. Number of cristae observed in mitochondrial sections by transmission electron microscopy.

Cell line	Mito section count	Number of cristae
29-13	239	231
KREPA6 KD tet+	227	213
BS	225	22

Figure 1






Figure 2

A



B



C




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7





Figure 8



Conclusions and perspectives

This thesis concerns two aspects of RNA editing in *Trypanosoma brucei*: 1) functional analysis of accessory proteins playing some role in this process; 2) understanding the phenomena of dyskinetoplasty. This section will discuss the conclusions and perspectives of each of these parts separately.

1. Functional analysis of mitochondrial RNA binding complex 1.

RNAi-silencing of the protein TbrGG1 results in a decrease in the steady state levels of the assayed edited RNAs without affecting those of gRNAs. The tandem affinity purification (TAP) of TbrGG1 led to the discovery of ~14 proteins that we provisionally named mitochondrial (mt) RNA binding complex 1 (MRB1), although its status as a *bona fide* complex remains in doubt. This notion is supported by the overlapping yet diverse MRB1 compositions reported by other groups identifying it with the same technique. Furthermore, TbrGG1 associates with this complex in a RNA-mediated manner, an indication that MRB1 may represent a group of smaller complexes and/or monomers.

Four subunits of MRB1, guide RNA associated proteins (GAPs) 1 and 2 as well as the predicted Nudix hydrolase and RNA helicase, were chosen for further analysis by reverse genetics. RNAi silencing of the GAPs and RNA helicase resulted in a downregulation of *trans*-gRNA mediated RNA editing due to a depletion of the minicircle encoded gRNAs, while *cis*-editing of *cox2* persisted. Furthermore, the GAPs appear to be interacting partners, are essential in the bloodstream stage and have an interesting, punctuate localization in the mitochondrion of the procyclic stage. In our study, Nudix hydrolase knockdowns exhibited a general destability of maxicircle encoded RNAs without affecting gRNAs. Thus, a subset of MRB1 appears to have a role in gRNA biogenesis. A single mt RNA polymerase seems to transcribe both maxicircle and minicircle transcripts, seemingly precluding such a role of the GAPs and RNA helicase in gRNA expression.

Because studies of the proteins comprising MRB1 is just starting, there are several directions for future research. The issue with most priority is the heterogeneity of data pertaining to MRB1. The diverse compositions of isolated MRB1 is striking, although they contain a significant overlap of identified subunits, such as the GAPs,

TbRGGs, Nudix hydrolase and RNA helicase. Work by Weng and colleagues (2008) state that these proteins are divided into four smaller complexes that are linked by RNA. The complexes are: 1) guide RNA binding complex (GRBC), which contains the RNA helicase and GAPs; 2) MERS, which contains the Nudix hydrolase; 3) the heterotetrameric MRB complex (Schumacher et al., 2006; Zíková et al., 2008); 4) KPAP complex, containing TbRGG1 (Etheridge et al., 2008). Data presented in this thesis, such as the variability of RNAi-silencing phenotypes, supports the notion that MRB1 represents smaller complexes. However, the division into these four complexes does not take into account all of the isolated MRB1 proteins or glaring absences of key proteins in other preparations, such as KPAP, MRB1 and 2 or the pentatricopeptide repeat proteins.

Clearly, a systematic approach is needed to define true protein-protein interactions between MRB1 proteins, such as a yeast-2-hybrid based approach, as utilized for defining interactions among 20S editosome subunits (Schnauffer et al., 2003). This effort in conjunction with the continued screening of other MRB1-subunit knockdowns for phenotypes such as gRNA depletion may allow for definition of smaller complexes and/or monomers in this collection of proteins.

The Nudix hydrolase phenotype described in this work counters that reported by Weng and co-workers (2008). However, our results are consistent with a role that these enzymes typically have in the repair of oxidative damage to nucleotides (McLennan, 2006). We verified that maxicircle DNA was not mutagenized in the Nudix-hydrolase knockdowns by sequencing several clones of PCR-amplified *cox1* (data not shown). To address whether this Nudix hydrolase has the capacity to repair oxidized ribonucleosides, it can be expressed in bacteria containing Nudix null-mutations to determine whether it rescues this repair pathway by measuring growth.

2. Understanding dyskinetoplasty

The dyskinetoplastic (dk) *Trypanosoma equipped* and akinetoplastic (ak) *Trypanosoma evansi* are important veterinarian pathogens that have a worldwide distribution despite exhibiting aberrant or complete loss of kDNA, respectively. Analysis of these trypanosomes employing molecular and classical-parasitology approaches indicated that they are strains of *T. brucei* and should be given subspecies status. The dk state seems to

arise spontaneously with the unfaithful replication minicircles. Although the kDNA network appears intact, it is only comprised of a few minicircle classes and a drastically diminished repertoire of gRNA genes. Since the selection pressure for maintaining maxicircles is no more, these molecules accumulate mutations such as deletions and are eventually lost. Eventually, the ak state is attained, in which all kDNA is lost. Because dk/ak cells prosper by fermentation of glucose, they can be considered to be petite mutants of *T. brucei*, which may arise repeatedly in an autonomous fashion in nature.

The existence of these trypanosomes ostensibly contradicts the finding that RNA editing is essential in the bloodstream stage (Schnauffer et al., 2001), presumably for the extensive editing of a transcript provisionally assigned to be F₀F₁ ATP synthase (complex V) subunit 6 (A6). This respiratory complex is essential in the bloodform to maintain membrane potential of the mitochondrion. Elegant studies have shown that compensatory mutations to another subunit in a laboratory-induced dk strain compensates for the absence of this mt encoded subunit in these cells (Schnauffer et al., 2005). We addressed the issue of whether this subunit is incorporated into complex V indirectly. By comparing the differential RNAi phenotypes of MRB1 knockdowns, affecting only mt encoded subunits of the electron transport chain, with KREPA6, which affects all respiratory chain complexes including complex V. We show the accumulation of the matrix F₁ moiety in the latter cells, while the assembly of the complex is not affected in the former. The oligomycin sensitivity of complex V ATPase activity is reduced in KREPA6-silenced cells as well. The accumulation F₀ is also observed in dk and ak cell lines as compared to *T. brucei* bloodforms. We conclude from these results that A6 is indeed incorporated into complex V. Furthermore, we propose that the F₁ moiety exists as a dimer.

The ongoing sequencing of the *T. evansi* ak strain used in these studies may identify candidate mutations that may compensate for the absence of A6 in these cell lines. Such mutations can be verified by expressing these in a RNA-editing incompetent cell line, such as the RNA ligase KREL1 knockouts (Schnauffer et al., 2005) or GAP knockdowns, to see whether growth is rescued.

References

Etheridge,R.D., Aphasizheva,I., Gershon,P.D., and Aphasizhev,R. (2008). 3' adenylation determines mRNA abundance and monitors completion of RNA editing in *T. brucei* mitochondria. *EMBO J.* 27, 1596-1608.

McClellan,A.G. (2006). The Nudix hydrolase superfamily. *Cell. Mol. Life Sci.* 63, 123-143.

Schnauffer,A., Panigrahi,A.K., Panicucci,B., Igo,R.P., Jr., Wirtz,E., Salavati,R., and Stuart,K. (2001). An RNA ligase essential for RNA editing and survival of the bloodstream form of *Trypanosoma brucei*. *Science* 291, 2159-2162.

Schnauffer,A., Clark-Walker,G.D., Steinberg,A.G., and Stuart,K. (2005) The F1-ATP synthase complex in bloodstream stage trypanosomes has an unusual and essential function. *EMBO J.* 24, 4029–4040.

Weng,J., Aphasizheva,I., Etheridge,R. D., Huang,L., Wang,X., Falick,A. M. and Aphasizhev,R. (2008). Guide RNA-binding complex from mitochondria of trypanosomatids. *Mol. Cell* 32, 1-12.